

Product Data Sheet

PE anti-mouse FR4 (Folate Receptor 4)

Catalog # / Size: 125107 / 25 µg

125108 / 100 µg

Clone: TH6

Isotype: Rat IgG2b, κ

Reactivity: Mouse Folate Receptor 4

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



BALB/c mouse splenocytes stained with TH6 PE and CD3 (145-2C11

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For

immunofluorescent staining, the suggested use of this reagent is ≤0.06 μg per million cells in 100 μl volume. It is

recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Yamaguchi T et al. 1007: Immunity 27(1)145

Description: Folate receptor 4 (FR4) is a surface receptor for folic acid (Vitamin B9). It was recently reported that FR4 is highly and

constitutively expressed on mouse CD4+ CD25+ natural regulatory T cells (Treg). The high expression of this surface marker on Treg can be used in combination with CD4 and CD25 to distinguish Treg from other types of T cells Since FR4 is a surface marker, this antibody may be used in combination with anti-CD4 and/or CD25 to sort living T reg cells

93

for functional studies.

Antigen References: 1. Yamaguchi T et al 2007. Immunity 27:145

2. Spiegelstein O et al. 2000 Gene 258:117

Related Products: Product Clone Application FC, ICC, ICFC Cell Staining Buffer PE Rat IgGŽb, κ Isotype Ctrl RTK4530 FC, ICFC RBC Lysis Buffer (10X) FC, ICFC FC TruStain fcX™ (anti-mouse CD16/32)

