

Product Data Sheet

Alexa Fluor® 647 anti-mouse/rat/human FOXP3 Flow Kit

Catalog # / Size: 320021 / 25 tests

Clone: 150D

Reactivity: Mouse, Rat, Human, **Cross-Reactivity:** Cynomolgus, Rhesus, Baboon

Storage: This kit is guaranteed for three months. Upon receipt, store at 4°C and protected from prolonged exposure to light. **Do not freeze.**


Applications:

Applications: ICFC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent intracellular staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 µl per million cells or 5 µl per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633 nm / 635 nm.

** Alexa Fluor® 647 is a registered trademark of Molecular Probes, Inc. Alexa Fluor® 647 dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.



C57BL/6 splenocytes surface stained with CD4 FITC, then intracellular stained with Alexa Fluor® 647 FOXP3 (150D) Flow Kit. Quadrant markers were set based on staining with Alexa Fluor® 647 mouse IgG1, κ isotype control

Application Notes: **Materials Provided:**

1. Alexa Fluor® 647 anti-mouse/rat/human FOXP3
25 tests
2. Alexa Fluor® 647 Mouse IgG1, κ isotype control
25 tests
3. FOXP3 Fix/Perm buffer set, 100 tests (Cat. No. 421403)

Materials not included:

Cell Staining Buffer (Cat. No. 420201)

Surface Staining & FOXP3 Buffer Preparation:

Centrifugation steps: perform at 250Xg for 5min

Incubation steps: perform at room temperature

1. Perform cell surface staining if necessary (See protocol: Cell Surface Immunofluorescence Staining Protocol).
2. Prepare 1X buffer solutions of FOXP3 Fix/Perm buffer and FOXP3 Perm buffer in PBS.

NOTE: The FOXP3 Perm buffer (10X) may have crystallization or precipitation observed when it is stored at 2-8°C, however, it is normal and does not affect the buffer performance. If there is a heavy precipitation observed after diluting to 1X working solution, it may be clarified by filtering.

Caution: The FOXP3 Fix/Perm buffer contains paraformaldehyde, which is toxigenic and mutagenic. Please handle with caution and wear gloves, lab coat and necessary protection to avoid direct body contact.

FOXP3 Intracellular Staining Procedures:

3. Add 1 ml of 1X BioLegend's FOXP3 Fix/Perm solution to each tube, resuspend the cells (gentle vortex) and incubate at room temperature in the dark for 20 minutes, then centrifuge and remove the supernatant. *The cell pellet will now be translucent and difficult to see; take care not to dislodge and accidentally aspirate cells at all later stages of staining protocol.*
4. Wash: resuspend cells in cell staining buffer (Cat. No. 420201); centrifuge, then discard the supernatant.
5. Wash: resuspend in 1ml 1X BioLegend's FOXP3 Perm buffer; centrifuge, then discard the supernatant.
6. Resuspend cells in 1ml 1X BioLegend's FOXP3 Perm buffer, incubate in the dark for 15 minutes; centrifuge, then discard the supernatant. Resuspend the pellet in 100 µl of 1X BioLegend's FOXP3 Perm buffer.
7. Add appropriate amount of fluorochrome conjugated anti-FOXP3 antibody and incubate at room temperature in the dark for 30 minutes.
8. Wash twice with cell staining buffer (see step 4) then resuspend in 0.5 ml cell staining buffer. Analyze with flow cytometer using appropriate instrument settings.

NOTE: BioLegend's FOXP3 Fix/Perm buffer set (Cat. No. 421403) is specifically developed and formulated for intracellular staining FOXP3 with minimum effect on surface fluorochrome staining and is highly recommended for



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

optimal result of FOXP3 intracellular immunofluorescence staining.

Application References:

1. Roncador G, *et al.* 2005 *Eur. J. Immunol.* 35:1681.
2. Mayack S, *et al.* 2006. *J. Immunol.* 176:2059. PubMed.
3. Yang ZZ, *et al.* 2006. *Blood* 107:3639.
4. Gavin MA, *et al.* 2006. *P. Natl. Acad. Sci. USA* 103:6659.
5. Groh V, *et al.* 2006. *Nature Immunology* 7:755.
6. Lewkowicz P, *et al.* 2006 *J. Immunol.* 177:7155.
7. Luke PPW, *et al.* 2006. *Amer. J. Transplant.* 6(9):2023.
8. Bamias G, *et al.* 2007. *J. Immunol.* 178:1809.
9. Valencia X, *et al.* 2007. *J. Immunol.* 178:2579. PubMed
10. Davidson TS, *et al.* 2007. *J. Immunol.* 178:4022.
11. MacDonald K PA, *et al.* 2007. *Blood* doi:10.1182/blood-2007-01-067249.
12. Jaffar Z, *et al.* 2007. *J. Immunol.* 179:6193.
13. Müller M, *et al.* 2007. *J. Immunol.* 179:2774.
14. Jordan JM, *et al.* 2008. *Infect Human.* 76:3717. PubMed
15. Golovina TN, *et al.* 2008. *J. Immunol.* 181:2855. PubMed
16. Fallarino F, *et al.* 2009. *J. Exp Med.* 206:2511. PubMed
17. Banham Alison, *et al.* 2009. *Vet Immunol and Immunop* 127.3-4:376-381
18. Klunker S, *et al.* 2009. *J. Exp Med.* PubMed
19. Haque A, *et al.* 2010. *J. Immunol.* 184:2583. PubMed

Description: FOXP3 is a 50 kD transcription factor, also known as Forkhead box protein P3, Scurfin, JM2, or IPEX. It is proposed to be a master regulatory gene and more specific marker of T regulatory cells than most cell surface markers (such as CD4 and CD25). Transduced expression of FOXP3 in CD4⁺/CD25⁻ cells has been shown to induce GITR, CD103, and CTLA4 and impart a T regulatory cell phenotype. FOXP3 is mutated in X-linked autoimmunity-allergic dysregulation syndrome (XLAAD or IPEX) in humans and in "scurfy" mice. Overexpression of FOXP3 has been shown to lead to a hypoactive immune state suggesting that this transcriptional factor is a central regulator of T cell activity. In human, unlike in mouse, two isoforms of FOXP3 have been reported: one (FOXP3) corresponding to the canonical full-length sequence; the other (FOXP3 $\delta 2$) lacking exon 2. The 150D monoclonal antibody reacts with human, mouse and rat FOXP3. The 150D antibody recognizes FOXP3 epitope encoded by exon 2.

BioLegend's Alexa Fluor® 647 anti-mouse/rat/human FOXP3 Flow Kit is designed and formulated specifically for immunofluorescence staining and flow cytometric analysis of mouse/rat/human FOXP3⁺ Treg cells in a mixed lymphocyte population. This kit is composed of Alexa Fluor® 647 conjugated anti-mouse/rat/human FOXP3 antibody and Alexa Fluor® 647 conjugated matching isotype control and the critical buffers. It is easy to use for identification of FOXP3⁺ Treg cells.

Antigen References:

1. Hori S, *et al.* 2003. *Science* 299:1057.
2. Fontenot JD, *et al.* 2003. *Nature Immunol.* 4:330.
3. Ferguson PJ, *et al.* 2000. *Am. J. Med. Genet.* 90:390.
4. Bennett CL, *et al.* 2001. *Nature Genet.* 27:20.
5. Allan SE, *et al.* 2005. *J. Clin. Invest.* 115:3276.

Related Products:

Product	Clone	Application
Cell Staining Buffer		FC, ICC, ICFC
Human Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD4 PE-Cy5/CD25 PE)	259D	FC, ICFC
Mouse Treg Flow™ Kit (FOXP3 Alexa Fluor® 488/CD4 APC/CD25 PE)	150D	FC, ICFC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.