

Product Data Sheet

Purified anti-mouse/human KLRG1 (MAFA)

Catalog # / Size: 138402 / 500 µg

Clone: 2F1/KLRG1

Isotype: Syrian hamster IgG

Immunogen: IL-2 activated NK cells from C57BL/6 mice

Reactivity: Mouse, Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.

Applications:

Applications: FC - Quality tested WB - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each


application.

Application Notes: Additional reported applications (for relevant formats) include: Western


Blotting¹.

This product may be used for research purposes only. It is not licensed for resale and may only be used by the buyer. This product may not be used and is not licensed for clinical assays, where the results of such assays are provided as a diagnostic service. If a diagnostic or therapeutic use is anticipated, then a license must be requested from the University of California. The availability of such diagnostic and therapeutic use license(s)


cannot be guaranteed from the University of California.



2. Robbins SH, *et al.* 2003. *J. Immunol.* 170:5876. (FC) 3. McMahon CW, *et al.* 2002. *J. Immunol.* 169:1444. (FC)



C57BL/6 mouse splenocytes stained with NK-1.1 (PK136) APC and purified 2F1/KLRG1 conjugated with



C57BL/6 mouse splenocytes stained with NK-1.1 (PK136) APC and Syrian hamster IgG PE isotype control

Description: Killer cell lectin-like receptor G1 (KLRG1) is the mouse homolog of the rat mast cell function-associated antigen (MAFA or 2F1-Ag). KLRG1 is a type II membrane glycoprotein that was first identified on the surface of rat mast cell line RBL-2H3. It is composed of a homodimer of glycosylated 30-38 kD subunits. Mouse and human homologs of KLRG1 are expressed by subsets of NK cells and lymphokine-activated killer (LAK) cells but not mast cells. KLRG1 is also expressed on subsets of CD8+ and CD4+ cells, including CD4+ and CD8+ effector/memory cells, potent regulatory CD4+ T cells. KLRG1 may be involved in regulating NK cell homeostasis. KLRG1 was recently found to

recognize cadherins and thus inhibit immune responses by regulating the effector function and the developmental

processes of NK and T cells.

Antigen References: 1. Grundemann C, et al. 2006. J. Immunol. 176:1311. 2. Blaser C, et al. 1998. J. Immunol. 161:6451. 3. Huntington ND, et al. 2007. J. Immunol. 178:4764. 4. Voehringer D, et al. 2001. J. Immunol. 167:4834.

Related Products: Product

Purified Syrian Hamster IgG Isotype Ctrl

Cell Staining Buffer RBC Lysis Buffer (10X) Clone SHG-1

Application FC, ICC, ICFC, IF, IHC, IP, WB FC, ICC, ICFC

FC, ICFC



