

FITC anti-mouse IgD

Catalog # / Size: 405703 / 50 µg
405704 / 500 µg

Clone: 11-26c.2a

Isotype: Rat IgG2a, κ


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 mouse splenocytes stained with 11-26c.2a FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: The 11-26c.2a antibody reacts with immunoglobulin D in all tested mouse haplotypes. The antibody binds membrane IgD expressed on most B cells. The 11-26c.2a antibody neither induces proliferation of splenic B cells nor induces B cell activation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections^{2,3}.

Application References:

1. Nitschke L, *et al.* 1993. *P. Natl. Acad. Sci. USA* 90:1887. (FC)
2. Weih D, *et al.* 2001. *J. Immunol.* 167:1909. (IHC)
3. Koni PA, *et al.* 2001. *J. Exp. Med.* 193:741. (IHC)
4. Ahuja A, *et al.* 2007. *J. Immunol.* 179:3351. (FC) PubMed
5. Haynes NM, *et al.* 2007. *J. Immunol.* 179:5099. (FC)
6. Good-Jacobson KL, *et al.* 2010. *Nat. Immunol.* 11:535. (FC) PubMed
7. Tomayko MM, *et al.* 2010. *J. Immunol.* 185:7146. PubMed
8. Park SY, *et al.* 2013. *J. Immunol.* 190:1094. PubMed

Description: Surface IgD is an important B cell differentiation marker.

Related Products:

Product
FITC anti-mouse/human CD45R/B220
Cell Staining Buffer
FITC anti-mouse CD19
FITC Rat IgG2a, κ Isotype Ctrl
FITC anti-mouse CD19

Clone
RA3-6B2

6D5
RTK2758
MB19-1

Application
FC
FC, ICC, ICFC
FC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.