

## Alexa Fluor® 647 anti-mouse Notch 1

**Catalog # / Size:** 130611 / 25 µg  
130612 / 100 µg

**Clone:** HMN1-12

**Isotype:** Armenian Hamster IgG

**Immunogen:** Notch1-Fc fusion protein


**Reactivity:** Mouse

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 647 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 647.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Mouse NOTCH-1 transfected cells stained with HMN1-12 Alexa Fluor® 647

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq 1.0$  µg per  $10^6$  cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

\*\* Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

**Application References:** 1. Moriyama Y, *et al.* 2008. *Int. Immunol.* 20:763.

**Description:** The Notch receptors are highly conserved from invertebrates to mammals. Notch 1 and Notch 2 exhibit the greatest structural similarity among the four mammalian Notch receptors. Notch 3 has a number of structural and functional differences. The binding of Notch 3 to its ligands results in the proteolysis of Notch and movement of intracellular portions of Notch into the nucleus. This translocation triggers a series of signaling processes. Notch 3 is primarily expressed in adult arterial smooth muscle cells. Notch 3 gene mutation can cause CADASIL, an inherited early stroke syndrome.

**Antigen References:** 1. Ehebauer ME, *et al.* 2006. *Biochem. J.* 392:13.  
2. Shimizu K, *et al.* 2000. *Mol. Cell. Biol.* 20:18.  
3. Tanigaki K, *et al.* 2007. *Nature Immunol.* 8:4514.  
4. Kraman M, *et al.* 2005. *FASEB J.* 19:1311.

### Related Products:

<b>Product</b>	<b>Clone</b>	<b>Application</b>
Alexa Fluor® 647 Armenian Hamster IgG Isotype Ctrl	HTK888	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
TruStain fcX™ (anti-mouse CD16/32)	93	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.