

Alexa Fluor® 647 anti-mouse/human CD15 (SSEA-1)

Catalog # / Size: 125607 / 25 tests
125608 / 100 tests

Clone: MC-480

Isotype: Mouse IgM, κ


Immunogen: Mouse F9 Teratocarcinoma Stem Cells (X-irradiated)

Reactivity: Mouse, **Cross-Reactivity:** Human

Preparation: The antibody was conjugated with Alexa Fluor® 647 under optimal conditions, and is at >85% purity. The solution is free of unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



F9 (mouse embryonic carcinoma cell line) stained with MC-480 Alexa Fluor® 647

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μ l per million cells or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation¹, Western blotting¹, and immunohistochemistry¹ of acetone-fixed frozen tissue sections and formalin-fixed paraffin-embedded sections.

Application References: 1. Solter D and Knowles BB. 1978. *Proc. Natl. Acad. Sci. USA.* 75:5565. (IHC, IP, WB)

Description: The MC-480 antibody reacts with mouse and human Stage-Specific Embryonic Antigen-1 (SSEA-1). It is a Lewis blood group related carbohydrate antigen, also known as X-hapten, Lewis X, 3-FAL, 3-fucosyl-N-acetylactosamine, or CD15. The expression pattern of SSEA-1 antigen is different in humans than in mice. In mice, SSEA-1 is expressed on embryonic stem cells (ES), embryonal carcinoma cells (EC), 8-cell to blastocyst embryos, and a subset of embryonic inner cell mass. The expression on murine ES cells is decreased upon differentiation. In humans, however, SSEA-1 is not found on undifferentiated ES cells, but its expression is upregulated along with differentiation. CD15 is highly expressed on adult human granulocytes. It has been reported that SSEA-1 plays a role in cell adhesion and regulation of cell differentiation.

Antigen References: 1. Solter D and Knowles BB. 1978. *Proc. Natl. Acad. Sci. USA.* 75:5565.
2. Harris JF, *et al.* 1984. *J. Immunol.* 132:2502.
3. Gooi HC, *et al.* 1981. *Nature* 292:156.
4. Cui L, *et al.* 2004. *J. Histochem. Cytochem.* 52:1447.

Related Products:

Product
Alexa Fluor® 647 Mouse IgM, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
MM-30

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.