

## PE/Cy7 anti-mouse Podoplanin

**Catalog # / Size:** 127411 / 25 µg  
127412 / 100 µg

**Clone:** 8.1.1

**Isotype:** Syrian Hamster IgG


**Reactivity:** Mouse Podoplanin

**Preparation:** The antibody was purified by affinity chromatography and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.2 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Mouse thymic epithelial stromal cell line TE-71 stained with 8.1.1 PE/Cy7

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** Additional reported applications (for the relevant formats) include: immunohistochemistry<sup>6</sup>.

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only.

**Application References:**

1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
2. Farr AG, *et al.* 1992. *J. Exp. Med.* 176:1477.
3. Bekiaris V, *et al.* 2008. *J. Immunol.* 180:6768.
4. Algars A, *et al.* 2011. *Blood* 117:4387. PubMed
5. Reis VO, *et al.* 2012. *Immunobiology.* 217:831. PubMed
6. Kaji C, *et al.* 2012. *Acta. Histochem. Cytochem.* 45:227. (IHC)

**Description:** The mucin-type glycoprotein podoplanin is thought to be involved in the development of the lymphatic vascular system. Podoplanin is named after its expression in the kidney glomerular epithelial cells (podocytes). It has a potential role in tumor progression.

**Antigen References:**

1. Farr A, *et al.* 1992. *J. Histochem. Cytochem.* 40:651.
2. Schacht V, *et al.* 2005. *Am. J. Pathol.* 166:913.

### Related Products:

**Product**  
Cell Staining Buffer  
RBC Lysis Buffer (10X)  
TruStain fcX™ (anti-mouse CD16/32)

### Clone

93

### Application

FC, ICC, ICFC  
FC, ICFC  
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.