


Anti-Mouse CD117 (c-Kit) PE-Cy7

Catalog Number: 25-1172

Also Known As: cKit, Steel Factor Receptor

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57BL/6 bone marrow cells with Anti-Human/Mouse CD45R (B220) FITC (cat. 11-0452) and 0.5 ug of Rat IgG2b kappa Isotype Control PE-Cy7 (cat. 25-4031) (left) or 0.5 ug of Anti-Mouse CD117 (c-Kit) PE-Cy7 (right). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD117 (c-Kit) PE-Cy7

REF Catalog Number: 25-1172

Clone: ACK2

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2b, kappa

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material. This Cy7 tandem dye is sensitive to photo-induced oxidation. Protect this vial from light during storage, handling & experimental procedures.



LOT Batch Code: Refer to Vial



Use By: Refer to Vial



Caution, contains Azide

Description

The ACK2 monoclonal antibody reacts with mouse CD117, also known as c-Kit receptor, Steel factor receptor and stem cell factor receptor. A member of the tyrosine kinase receptor family, this 145 kDa molecule is expressed by a majority of hematopoietic progenitor cells characterized in the mouse bone marrow as a small subset of cells positive for Sca-1 and Thy1 (Thy1^{lo}) and negative for lineage markers. The interaction of the mouse c-kit receptor and steel factor promotes the proliferation and differentiation of hematopoietic progenitor cells. CD117 is also expressed by mast cells and plays a role in signaling and activation of these cells. ACK2 has been reported to be a blocking antibody.

Applications Reported

The ACK2 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This ACK2 antibody has been tested by flow cytometric analysis of mouse bone marrow cell suspensions. This can be used at less than or equal to 1 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100 uL cell sample + 100 uL IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency/compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

References

Rico-Vargas, S., et al. 1994. c-kit expression by B cell precursors in mouse bone marrow. J. Immunol. 152: 2845.

Ito M, Kawa Y, et al. 1999. Removal of stem cell factor or addition of monoclonal anti-c-KIT antibody induces apoptosis in murine melanocyte precursors. J Invest Dermatol. 112: 796-801.

Vincent S, Segretain D, et al. 1998. Stage-specific expression of the Kit receptor and its ligand (KL) during male gametogenesis in the mouse: a Kit-KL interaction critical for meiosis. Development. 125: 4585-93.

Sato D, Lai ZF, et al. 1996. Impairment of Kit-dependent development of interstitial cells alters contractile responses of murine intestinal tract.

Am J Physiol. 271: G762-71.

Yoshida H, Kunisada T, et al. 1996. Distinct stages of melanocyte differentiation revealed by analysis of nonuniform pigmentation patterns. *Development*. 122: 1207-14.

Yoshinaga K, Nishikawa S, et al. 1991. Role of c-kit in mouse spermatogenesis: identification of spermatogonia as a specific site of c-kit expression and function. *Development*. 113:689-99.

Torihashi S, Ward SM, et al. 1995. c-kit-dependent development of interstitial cells and electrical activity in the murine gastrointestinal tract. *Cell Tissue Res*. 280:97-111.

Feng H, Sandlow JI, Sandra A. 1997. Expression and function of the c-kit proto-oncogene protein in mouse sperm. *Biol Reprod*. 57:194-203.

Feng H, Sandlow JI, Sandra A. 1998. The c-kit receptor and its possible signaling transduction pathway in mouse spermatozoa. *Mol Reprod Dev*. 49:317-26.

Prasanth SG and Ali S. 2003. Expression of proto-oncogene c-kit receptor in rats (*Rattus norvegicus*) and identification of a mutant mRNA transcript implicated in spermatogenic failure. *DNA Cell Biol*. 2003 Jul;22(7):447-56. (ACK2, rat cross-reactivity, PubMed)

Ismail RS, Okawara Y, et al. 1996. Hormonal regulation of the ligand for c-kit in the rat ovary and its effects on spontaneous oocyte meiotic maturation. *Mol Reprod Dev*. 1996 Apr;43(4):458-69. (ACK2, rat cross-reactivity and blocking, PubMed)

Kim HM, Shin HY, and Lee EH. 1998. Morphological alterations in rat peritoneal mast cells by stem cell factor. *Immunology*. 1998 Jun;94(2):242-6. (ACK2, rat cross-reactivity and blocking, PubMed)

Czechowicz A, Kraft D, Weissman IL, and Bhattacharya D. 2007. Efficient transplantation via antibody-based clearance of hematopoietic stem cell niches. *Science*. 23;318(5854):1296-9. (ACK2, in vivo depletion HSC, PubMed)

Related Products

11-0452 Anti-Human/Mouse CD45R (B220) FITC (RA3-6B2)

25-4031 Rat IgG2b K Isotype Control PE-Cy7

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486, 5,569,587 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com