

Product Data Sheet

PE anti-mouse TCR V γ 2

Catalog # / Size: 137705 / 50 μ g
137706 / 200 μ g

Clone: UC3-10A6

Isotype: Armenian hamster IgG

Immunogen: G8 mouse T cells


Reactivity: Mouse

Preparation: The antibody was purified by affinity chromatography and conjugated with PE under optimal conditions. The solution is free of unconjugated PE and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.2 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



C57BL/6 mouse lymphocytes stained with TCR V γ 2 (UC3-10A6) PE and CD3 (145-2C11) APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.125 μ g per million cells in 100 μ l volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

1. Dent AL, *et al.* 1990. *Nature* 343:714.
2. Kelly KA, *et al.* 1993. *Int. Immunol.* 5:331.
3. Sperling AI, *et al.* 1992. *J. Immunol.* 149:3200.
4. Sperling AI, *et al.* 1997. *J. Immunol.* 159:86.

Description: T cell receptor (TCR) V γ 2 bearing T lymphocytes make up a significant proportion of $\gamma\delta$ TCR cells in late fetal and adult peripheral lymphoid tissues. TCR $\gamma\delta$ T cells may play a role in immunological surveillance for stress-induced self-antigens. The frequency of V γ 2 expression in different strains varied from 12% to 54% in the TCR $\gamma\delta$ repertoire. Variations in the levels of V γ 2⁺ cells are not associated with MHC haplotype. High V γ 2 expression is influenced by the TCR- δ locus. Expanding V γ 2⁺ TCR $\gamma\delta$ cells in B6 mice overwhelmingly use a V δ 7⁺ δ chain except in the DBA/2 strain.

Antigen References:

1. Allison JP, *et al.* 1991. *Annu. Rev. Immunol.* 9:679.
2. O'Brien RL, *et al.* 2000. *J. Immunol.* 165:6472.

Related Products:

Product
PE Armenian Hamster IgG Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)
TruStain fcX™ (anti-mouse CD16/32)

Clone
HTK888

93

Application
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biollegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.