

Applications: ICFC- Quality tested

Product Data Sheet

FITC anti-human/mouse ZAP-70

| Catalog # / Size: | 313405 / 25 tests 313406 / 100 tests | |
|-------------------|---|---|
| Clone: | 1E7.2 | |
| Isotype: | Mouse IgG1 | |
| Immunogen: | GST-fusion protein corresponding to residues 282-307 of human ZAP-70 | |
| Reactivity: | Human, Mouse | |
| Preparation: | The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC. | 1 |
| Formulation: | Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA). | |
| Storage: | The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze. | |
| | | |

Applications:

Human peripheral blood lymphocytes intracellularly stained with 1E7.2 FITC and CD3 APC

Recommended Usage: Each lot of this antibody is quality control tested by intracellular immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 µl to 5 µl per test**. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation², Western blotting².

This product is for in vitro research use only. It is not to be used for commercial purposes. Use of this product to produce products for sale or for diagnostic, therapeutic or drug discovery purposes is prohibited. In order to obtain a license to use this product for commercial purposes, contact The Regents of the University of California.

Application References: 1. Qian D, et al. 1997. J. Exp. Med. 185:1253. (ICFC) 2. Crespo M, et al. 2003. New Engl. J. Med. 348:1764. (WB IP) 3. Jamroziak K, et al. 2009. Cancer Epidemiol Biomarkers Prev. PubMed

Description: ZAP-70 is a 70 kD protein tyrosine kinase associated with the zeta chain of the T cell receptor. It is expressed in T cells and NK cells and has been shown to be involved in T cell signaling. Defects in ZAP-70 have been linked to selective T cell defects. The ZAP-70 kinase undergoes multiple phosphorylation events after T cell receptor engagement and interacts with a number of proteins involved in signal transduction. Recently, ZAP-70 has been identified as an important prognostic marker in B-cell chronic lymphocytic leukemia (B-CLL). The 1E7.2 monoclonal antibody recognizes human and mouse ZAP-70 and has been shown to be useful for flow cytometry, Western blotting, and immunoprecipitation.

Antigen References: 1. Chan AC, *et al.* 1992. *Cell* 71:649. 2. Chan AC, *et al.* 1994. *Science* 264:1559. 3. Arpaia E, *et al.* 1994. *Cell* 76:947.

 Related Products: Product

 Cell Staining Buffer

 RBC Lysis Buffer (10X)

 Permeabilization Wash Buffer (10X)

 FITC Mouse IgG1, κ Isotype Ctrl (FC)

 PE Goat anti-mouse IgG (minimal x-reactivity)

 Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone

MOPC-21

Poly4053

Application FC, ICC, ICFC FC, ICFC ICC, ICFC, IHC FC FC FC, ICC, ICFC

!

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.