


Anti-Mouse CD199 (CCR9) PE-Cyanine7

Catalog Number: 25-1991

Also known as: C-C chemokine receptor type 9

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of BALB/c thymocytes with Anti-Mouse CD4 eFluor[®] 450 (cat. 48-0042), Anti-Mouse CD8a APC (cat. 17-0081) and 0.25 ug of Mouse IgG2a K Isotype Control PE-Cyanine7 (cat. 25-4724) (left) or 0.25 ug of Anti-Mouse CD199 (CCR9) PE-Cyanine7 (right). Samples were gated on CD4 single positive (orange histogram), CD8 single positive (blue histogram), or CD4+CD8+ double positive (purple histogram) cells.

Product Information

Contents: Anti-Mouse CD199 (CCR9) PE-Cyanine7

REF **Catalog Number:** 25-1991

Clone: eBioCW-1.2 (CW-1.2)

Concentration: 0.2 mg/mL

Host/Isotype: Mouse IgG2a

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light-sensitive material. This tandem dye is sensitive to photo-induced oxidation. Protect this vial from light during storage, handling & experimental procedures.

Batch Code: Refer to vial

Use By: Refer to vial

Contains sodium azide

Description

The eBioCW-1.2 monoclonal antibody reacts with mouse CCR9 (CD199), which is the receptor for thymus-expressed chemokine (TECK). CCR9 is a member of the G protein coupled receptor (GPCR) supergene family, and is involved in trafficking of T cell progenitors within the thymus. CCR9 expression during thymocyte development commences at the double-negative (DN) 3 stage (CD4-CD8-CD25+CD44-), peaks in the double-positive (DP) stage (CD4+CD8+CD25-CD44-), and is down-regulated in committed CD4+ or CD8+ single-positive (SP) thymocytes. CCR9-deficient mice show a mild impairment in thymocyte development. In the periphery, CCR9 is thought to be expressed on naïve CD8+ T cells, but not on naïve CD4+ T cells.

Applications Reported

This eBioCW-1.2 (CW-1.2) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBioCW-1.2 (CW-1.2) antibody has been tested by flow cytometric analysis of mouse thymocytes. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

Light sensitivity: This tandem dye is sensitive photo-induced oxidation. Please protect this vial and stained samples from light.

Fixation: Samples can be stored in IC Fixation Buffer (cat. 00-8222) (100 uL cell sample + 100 uL IC Fixation Buffer) or 1-step Fix/Lyse Solution (cat. 00-5333) for up to 3 days in the dark at 4°C with minimal impact on brightness and FRET efficiency/compensation. Some generalizations regarding fluorophore performance after fixation can be made, but clone specific performance should be determined empirically.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com

Anti-Mouse CD199 (CCR9) PE-Cyanine7

Catalog Number: 25-1991

Also known as: C-C chemokine receptor type 9

RUO: For Research Use Only. Not for use in diagnostic procedures.

References

Liu C, Saito F, Liu Z, Lei Y, Uehara S, Love P, Lipp M, Kondo S, Manley N, Takahama Y. Coordination between CCR7- and CCR9-mediated chemokine signals in pre-vascular fetal thymus colonization. *Blood*. 2006 Jun 29.

Wurbel MA, Malissen B, Campbell JJ. Complex regulation of CCR9 at multiple discrete stages of T cell development. *Eur J Immunol*. 2006 Jan;36(1):73-81. (CW-1.2, FC, Development of mAb, PubMed)

Zaballos A, Gutierrez J, Varona R, Ardavin C, Marquez G. Cutting edge: identification of the orphan chemokine receptor GPR-9-6 as CCR9, the receptor for the chemokine TECK. *J Immunol*. 1999 May 15;162(10):5671-5.

Related Products

17-0081 Anti-Mouse CD8a APC (53-6.7)

25-4724 Mouse IgG2a K Isotype Control PE-Cyanine7

48-0042 Anti-Mouse CD4 eFluor[®] 450 (RM4-5)

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486, 5,569,587 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com •
info@ebioscience.com