

Alexa Fluor® 488 anti-rat CD11b/c

Catalog # / Size: 201812 / 100 µg

Clone: OX-42

Isotype: Mouse IgG2a, κ

Immunogen: Rat peritoneal macrophages

Reactivity: Rat

Preparation: The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 488 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 488.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Applications:

Applications: FC - *Quality tested*
IHC, IF - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.

** Alexa Fluor® 488 is a registered trademark of Molecular Probes, Inc. Alexa Fluor® 488 dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemistry of acetone-fixed frozen sections^{1,2}, immunoprecipitation³, *in vivo* and *in vitro* blocking of C3bi binding^{3,4}.

Application References:

- Whiteland JL, et al. 1995. *J. Histochem. Cytochem.* 43:313. (IHC)
- Milligan CE, et al. 1991. *J. Comp. Neurol.* 314:125. (IHC)
- Robinson AP, et al. 1986. *Immunology* 57:239. (Block)
- Issekutz SE, et al. 1992. *Immunology* 76:655. (Block)
- Muehlbauer SM, et al. *Am. J. Pathol.* 177:735. (FC) PubMed

Description: The OX-42 antibody reacts with the CR3 complement (C3bi) receptor expressed on monocytes, granulocytes, macrophages, dendritic cells, NK cells, and a subset of lymphocytes. This antibody appears to recognize a common epitope shared between CD11b and CD11c (integrin α_M and α_X chains). The OX-42 antibody precipitates three polypeptides with apparent molecular weights of 160, 103, and 95 kD, respectively. This antibody has been shown to block the formation of complement-mediated rosettes and leukocyte migration.

Antigen References:

- Robinson AP, et al. 1986. *Immunology* 57:239.
- Barcaly AN. 1981 *Immunology* 42:593.

Related Products: Product

Cell Staining Buffer

Alexa Fluor® 488 Mouse IgG2a, κ Isotype Ctrl

Clone

MOPC-173

Application

FC, ICC, ICFC

FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.