

FITC anti-rat CD161

Catalog # / Size: 203105 / 100 µg

Clone: 10/78

Isotype: Mouse IgG1, κ

Immunogen: LEW rat splenic NK cells


Reactivity: Rat

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



LOU rat splenocytes stained with 10/78 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation³ and immunohistochemical staining of acetone-fixed frozen sections^{1,2}. Clone 10/78 is not suitable for immunohistochemical staining of formalin-fixed paraffin-embedded sections.

Application References:

1. Sedgwick JD, *et al.* 1998. *J. Immunol.* 160:5320. (IHC)
2. Tliba O, *et al.* 2002. *Vet. Res.* 33:327. (IHC)
3. Kraus E, *et al.* 1996. *Eur. J. Immunol.* 26:2582. (IP)
4. Treacy O, *et al.* 2012. *PLoS One.* 7:e42662. PubMed

Description: CD161 molecules, known as NKR-P1, are a family of about 30 kD type II transmembrane C-type lectin-like receptors and are expressed on the cell membrane as disulphide-linked homodimer. Rat NKR-P1 receptors are primarily expressed on NK cells, a subset of T cells, dendritic cells, and activated monocytes. Carbohydrate antigens with GalNac and GlcNac moieties are the ligands for NKR-P1 molecules. CD161 receptors are thought to be involved in the regulation of NK and NKT cell function. Three rat NKR-P1 genes have been described, NKR-P1A, NKR-P1B, NKR-P1B* (or NKR-P1D). 10/78, similar like 3.2.3 antibody, recognizes a common epitope of NKR-1A (CD161a) and NKR-P1B (CD161b). NKR-P1A does not contain ITIM structure and is an activating receptor, while NKR-P1B contains an ITIM and displays an inhibitory function.

Antigen References:

1. Ryan J, *et al.* 1991. *J. Immunol.* 147:3244.
2. Chambers WH, *et al.* 1989. *J. Exp. Med.* 169:1373.
3. Pospisil M, *et al.* 2000. *Int. J. Oncol.* 16(2):267.
4. Scriba A, *et al.* 1997. *J. Leukoc. Biol.* 62(6):741.
5. Brissette-Storkus CS, *et al.* 2002. *J. Leukoc. Biol.* 71(6):941.
6. Li J, *et al.* 2003. *Int. Immunol.* 15(3):411.

Related Products:

Product
Cell Staining Buffer

FITC Mouse IgG1, κ Isotype Ctrl (FC)

Clone

MOPC-21

Application

FC, ICC, ICFC
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.