

Purified anti-rat CD53

Catalog # / Size: 204302 / 500 µg

Clone: OX-44

Isotype: Mouse IgG1, κ

Immunogen: Rat T cell blasts


Reactivity: Rat

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



LOU rat splenocytes stained with purified OX-44, followed by anti-mouse IgG PE.

Applications:

Applications: FC - *Quality tested*
IHC , IP, IF, Cell Adhesion - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.25 µg per 10^6 cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, immunofluorescence microscopy, immunohistochemical staining for formalin-fixed paraffin-embedded tissue sections, and induction of cell adhesion.

Application References: 1. Lazo PA, *et al.* 1997. *Cellular Immunol.* 178:132

Description: CD53 is a 35-45 kD type III tetraspan membrane protein. It is expressed on all leukocytes and a subset of thymocytes, but not on platelets, erythrocytes, or non-hematopoietic cells. CD53 has been proposed to be involved in signal transduction and thymopoiesis. It has been shown to interact with a number of proteins, including IL-4, CD20, CD2, CD9, IL-2, VLA-4, and CD4. Cross-linking of CD53 promotes B cell activation. The CD53 antigen is highly glycosylated and treatment with endoglycosidase F reduces the apparent molecular weight of this protein by approximately 25kD.

Antigen References: 1. Lazo PA, *et al.* 1997. *Cellular Immunol.* 178:132

Related Products:

Product
Purified Mouse IgG1, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
MOPC-21

Application
FC, ICFC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.