

## Purified anti-human CD4

**Catalog # / Size:** 317401 / 25 µg  
317402 / 100 µg

**Clone:** OKT4

**Isotype:** Mouse IgG2b, κ


**Reactivity:** Human, **Cross-Reactivity:** Chimpanzee, Cynomolgus, Rhesus

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified OKT4, followed by anti-mouse IgGs FITC

## Applications:

**Applications:** FC - *Quality tested*  
IHC - *Reported in the literature*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.5 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

**Application Notes:** The OKT4 antibody binds to the D3 domain of CD4 and does not block HIV binding. Additional reported applications (for the relevant formats) include: immunohistochemistry of frozen sections and blocking of T cell activation. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 317404).

**Application References:**

- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Reinherz EL, *et al.* 1979. *Proc. Natl. Acad. Sci.* 76:4061.
- Kmieciak M, *et al.* 2009. *J. Transl. Med.* 7:89. (FC) PubMed
- Cicin-Sain L, *et al.* 2010. *J. Immunol.* 184:6739. PubMed
- Rosenzweig M, *et al.* 2001. *J. Med. Primatol.* 30:36.
- Linder J, *et al.* 1987. *Am. J. Pathol.* 127:1.
- Boche D, *et al.* 1999. *J. Neurovirol.* 5:232. (IHC)

**Description:** CD4, also known as T4, is a 55 kD single-chain type I transmembrane glycoprotein expressed on most thymocytes, a subset of T cells, and monocytes/macrophages. CD4, a member of the Ig superfamily, recognizes antigens associated with MHC class II molecules and participates in cell-cell interactions, thymic differentiation, and signal transduction. CD4 acts as a primary receptor for HIV, binding to HIV gp120. CD4 has also been shown to interact with IL-16.

**Antigen References:**

- Center D, *et al.* 1996. *Immunol. Today* 17:476.
- Gaubin M, *et al.* 1996. *Eur. J. Clin. Chem. Clin. Biochem.* 34:723.

**Related Products:**

**Product**  
Purified Mouse IgG2b, κ Isotype Ctrl  
Cell Staining Buffer  
RBC Lysis Buffer (10X)

**Clone**  
MPC-11

**Application**  
FC, ICFC, ICC, IF, IHC, IP, WB  
FC, ICC, ICFC  
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.