


## Anti-Human CD51/CD61 (Integrin alpha v beta 3) Functional Grade Purified

Catalog Number: 16-0519

Also Known As: Integrin  $\alpha$ v $\beta$ 3, vitronectin Receptor

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of platelets with 0.5  $\mu$ g of Mouse IgG1 kappa Isotype Control Purified (cat.14-4714) (open histogram), or 0.5  $\mu$ g of Anti-Human CD51/CD61 (Integrin alpha V beta 3) Purified followed by Anti-Mouse IgG FITC (cat. 11-4011) (filled histogram).

### Product Information

**Contents:** Anti-Human CD51/CD61 (Integrin alpha v beta 3) Functional Grade Purified

 **Catalog Number:** 16-0519

**Clone:** 23C6

**Concentration:** 1 mg/mL


**Host/Isotype:** Mouse IgG1, kappa

**HLDA Workshop:** IV P18

**Handling Conditions:** Use in sterile environment.

**Endotoxin Level:** Less than 0.001 ng/ $\mu$ g antibody, as determined by the LAL assay.

**Formulation:** aqueous buffer, no sodium azide

 **Temperature Limitation:** Store at 2-8°C.

 **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

### Description

The 23C6 monoclonal antibody reacts with the human CD51/CD61 dimer, also known as the integrin  $\alpha$ v $\beta$ 3. CD51, an ~120 kDa surface molecule can also non-covalently associate with other  $\beta$  subunits of the integrin family including  $\beta$ <sub>1</sub> (CD29),  $\beta$ <sub>5</sub> and  $\beta$ <sub>6</sub> to form receptors for extracellular matrix components. Heterodimers of CD51/CD61 are expressed by melanoma cells, endothelial cells and osteoclasts and at very low levels by platelets. The CD51/CD61 complex mediates adhesion to fibrinogen, fibronectin, vitronectin and thrombospondin.

### Applications Reported

This 23C6 antibody has been reported for use in flow cytometric analysis, and immunohistology staining of frozen tissue sections. 23C6 has also been reported in blocking of some adhesive processes.

### Applications Tested

This 23C6 antibody has been tested by flow cytometric analysis of human melanoma cell line and peripheral blood. This can be used at less than or equal to 1  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

### References

Schlossman, S., L. Bloumsell, et al. eds (1995). Leucocyte Typing V: White Cell Differentiation Antigens. Oxford University Press. New York.

### Related Products

11-4011 Anti-Mouse IgG FITC

14-4714 Mouse IgG1 K Isotype Control Purified (P3.6.2.1)

---

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • [www.eBioscience.com](http://www.eBioscience.com) • [info@eBioscience.com](mailto:info@eBioscience.com)