

Product Data Sheet

PerCP/Cy5.5 anti-human CD5

Catalog # / Size: 300619 / 25 tests

300620 / 100 tests

Clone: UCHT2

Isotype: Mouse IgG1, κ

Workshop Number: III 518

Reactivity: Human, Cross-Reactivity: Chimpanzee, Capuchin Monkey, Common

Preparation: The antibody was purified by affinity chromatography, and conjugated with PerCP/Cy5.5 under optimal conditions. The solution is free of unconjugated


PerCP/Cy5.5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes were stained with CD5 (ÚCHT2) PerCP/Cy5.5 (filled histogram) or mouse IgG1, k PerCP/Cy5.5 (open histogram).

FC, ICC, ICFC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μ l per million cells or 5 μ l per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

* PerCP/Cy5.5 has a maximum absorption of 482 nm and 564 nm and a

maximum emission of 690 nm.

Application Notes: Additional reported applications (for the relevant formats) include: Western blotting² and immunohistochemical

staining of acetone-fixed frozen sections^{2,5}.

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed

for research use only.

Application References: 1. Knapp W, et al. 1989. Leucocyte Typing IV Oxford University Press. New York.

Renaudineau Y, et al. 2005. Blood 106:2781. (WB IHC)
 Porter JC and Hogg N. 1997. J. Cell Biol. 138:1437.
 Saliba AE, et al. 2010. P. Natl. Acad. Sci. USA 107:14524. PubMed
 Kap Y, et al. 2009. J. Histochem. Cytochem. 57:1159. (IHC)

Description: CD5 is a 67 kD single chain type I glycoprotein also known as Leu-1, Ly-1 and T1. It is a member of the scavenger

receptor superfamily found on T cells, thymocytes, B cell subsets, chronic B lymphocytic leukemia (B-Cells), and peripheral blood dendritic cells. CD5 modulates T and B cell receptor signaling, thymocyte maturation, and T-B cell

interactions upon binding to ligands such as CD72.

Antigen References: 1. Kipps T. 1988. Adv. Immunol. 47:117.

2. Resnick D, et al. 1993. Trends Biochem. Sci. 19:5.

3. Wood GS, et al. 1992. Am. J. Pathol. 14:789.

Application Related Products: Product Clone

FC, ICC, ICFC FC, ICFC Cell Staining Buffer RBC Lysis Buffer (10X)
PerCP/Cy5.5 Mouse IgG1, κ Isotype Ctrl
Human TruStain FcX™ (Fc Receptor Blocking Solution) MOPC-21 FC, ICFC



