


Anti-Mouse CD54 (ICAM-1) Functional Grade Purified

Catalog Number: 16-0541

Also Known As: Intercellular adhesion molecule-1, ICAM1

RUO: For Research Use Only



Staining of mouse splenocytes with Anti-Mouse CD54 (ICAM-1) FITC (left) or PE (right). Autofluorescence is shown via open histogram. Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD54 (ICAM-1) Functional Grade Purified

REF **Catalog Number:** 16-0541

Clone: YN1/1.7.4

Concentration: 1 mg/ml

Host/Isotype: Rat IgG2b, κ


Handling Conditions: Use in sterile environment.

Endotoxin Level: Less than 0.001 ng/ μ g antibody, as determined by the LAL assay.

Formulation: aqueous buffer, no sodium azide

 **Temperature Limitation:** Store at 2-8°C.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

Description

The YN1/1.7.4 monoclonal antibody reacts with mouse CD54 (InterCellular Adhesion Molecule-1, ICAM-1), a 90-110 kDa transmembrane glycoprotein expressed by monocytes, lymphocytes, dendritic cells, and endothelial cells. Expression of CD54 is upregulated on activated lymphocytes. Interaction of CD54 with its ligands CD11a and CD11b is important in the inflammatory response. The YN1/1.7.4 monoclonal antibody recognizes a different epitope than the eBioKAT-1 monoclonal antibody.

Applications Reported

The YN1/1.7.4 antibody has been reported for use in flow cytometric analysis. YN1/1.7.4 has also been reported in blocking of CD54 in functional studies.

Applications Tested

The YN1/1.7.4 antibody has been tested by flow cytometric analysis of mouse splenocyte suspensions. This can be used at less than or equal to 0.5 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10^5 to 10^8 cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

- Burns, A. R., F. Takei, et al. (1994). "Quantitation of ICAM-1 expression in mouse lung during pneumonia." *J Immunol* 153(7): 3189-98.
- Fabry, Z., M. M. Waldschmidt, et al. (1992). "Adhesion molecules on murine brain microvascular endothelial cells: expression and regulation of ICAM-1 and Lgp 55." *J Neuroimmunol* 36(1): 1-11.
- Jevnikar, A. M., R. P. Wuthrich, et al. (1990). "Differing regulation and function of ICAM-1 and class II antigens on renal tubular cells." *Kidney Int* 38(3): 417-25.
- Kumasaka, T., W. M. Quinlan, et al. (1996). "Role of the intercellular adhesion molecule-1(ICAM-1) in endotoxin-induced pneumonia evaluated using ICAM-1 antisense oligonucleotides, anti-ICAM-1 monoclonal antibodies, and ICAM-1 mutant mice." *J Clin Invest* 97(10): 2362-9.

Related Products

- 11-4317 Streptavidin FITC
- 11-4811 Anti-Rat IgG FITC
- 12-4317 Streptavidin PE
- 13-4813 Anti-Rat IgG Biotin (Polyclonal)

16-4031 Rat IgG2b K Isotype Control Functional Grade Purified
17-4317 Streptavidin APC

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com