


Anti-Human CD244 Functional Grade Purified

Catalog Number: 16-2449

Also Known As: 2B4, p38, SLAMF4

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of normal human peripheral blood cells with Anti-Human CD8a APC (cat. 17-0088) and 0.25 ug of Mouse IgG1 K Isotype Control Functional Grade Purified (cat. 16-4714) (left) or 0.25 ug of Anti-Human CD244 Functional Grade Purified (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Human CD244 Functional Grade Purified

REF **Catalog Number:** 16-2449

Clone: eBioPP35 (PP35)

Concentration: 1 mg/mL

Host/Isotype: Mouse IgG1


Handling Conditions: Use in sterile environment.

Endotoxin Level: Less than 0.001 ng/ug antibody, as determined by the LAL assay.

Formulation: aqueous buffer, no sodium azide

 **Temperature Limitation:** Store at 2-8°C.

LOT **Batch Code:** Refer to Vial

 **Use By:** Refer to Vial

Description

The eBioPP35 monoclonal antibody reacts with human CD244, also known as 2B4, p38, NK-cell activation-inducing ligand, signaling lymphocyte activation molecule family member 4, and natural killer cell receptor 2B4. Human CD244 is a 38 kD protein expressed on NK cells, a subset of CD8+ T cells, $\gamma\delta$ T cells, monocytes, basophils, mast cells, and eosinophils. Binding of CD244 to its ligand, CD48, results in NK cell activation. Costaining experiments with eBioC1.7 suggest that both antibodies recognize different epitopes.

Applications Tested

This eBioPP35 (PP35) antibody has been tested by flow cytometric analysis of human peripheral blood cells.

References

Moretta A, Bottino C, Tripodi G, Vitale M, Pende D, Morelli L, Augugliaro R, Barbaresi M, Ciccone E, Millo R, et al. Novel surface molecules involved in human NK cell activation and triggering of the lytic machinery. *Int J Cancer Suppl.* 1992;7:6-10. (**PP35**, mAb development, FA, PubMed)

Parolini S, Bottino C, Falco M, Augugliaro R, Giliani S, Franceschini R, Ochs HD, Wolf H, Bonnefoy JY, Biassoni R, Moretta L, Notarangelo LD, Moretta A. X-linked lymphoproliferative disease. 2B4 molecules displaying inhibitory rather than activating function are responsible for the inability of natural killer cells to kill Epstein-Barr virus-infected cells. *J Exp Med.* 2000 Aug 7;192(3):337-46. (**PP35**, FC, FA, PubMed)

Sivori S, Falco M, Marcenaro E, Parolini S, Biassoni R, Bottino C, Moretta L, Moretta A. Early expression of triggering receptors and regulatory role of 2B4 in human natural killer cell precursors undergoing in vitro differentiation. *Proc Natl Acad Sci U S A.* 2002 Apr 2;99(7):4526-31. (**PP35**, FC, PubMed)

Related Products

16-4714 Mouse IgG1 K Isotype Control Functional Grade Purified (P3.6.2.1)

16-5838 Anti-Human CD244 Functional Grade Purified (eBioC1.7 (C1.7))

17-0088 Anti-Human CD8a APC (RPA-T8)

