

Purified anti-human CD11a

Catalog # / Size: 301202 / 100 µg

Clone: HI111

Isotype: Mouse IgG1, κ

Workshop Number: IV N231


Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Capuchin Monkey, Horse (Equine), Cattle (Bovine, Cow), Sheep (Ovine), Dog (Canine), Rabbit (Lapine)

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified HI111, followed by anti-mouse IgGs FITC

Applications:

Applications: FC - *Quality tested*
 IHC, WB - *Reported in the literature*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 0.5 µg per 10⁶ cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen sections, Western blotting², and blocking of cell-cell interaction and inhibition the binding of ICAM-1⁴. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 301214).

Application References:

- Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press New York.
- Leite F, *et al.* 2002. *Infect. Immun.* 70:4336.
- Jiang Y, *et al.* 2005. *Clin. Hemorheol. Microcircul.* 32:261.
- Bécharde D, *et al.* 2001. *J. Immunol.* 167:3099.
- Sithu SD, *et al.* 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
- Choi EY, *et al.* 2008. *Blood* 111:3607. PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)

Description: CD11a is a 170-180 kD type I transmembrane glycoprotein also known as LFA-1α chain and integrin α_L subunit. CD11a non-covalently associates with integrin β₂ (CD18) to form LFA-1. It is expressed on all leukocytes, including B and T lymphocytes, monocytes, macrophages, neutrophils, basophils and eosinophils. It is absent on non-hematopoietic tissues and platelets. CD11a plays a central role in leukocyte cell-cell interactions and is important in lymphocyte costimulation. CD11a/CD18 binds to ICAM-1 (CD54), ICAM-2 (CD102), and ICAM-3 (CD50).

Antigen References:

- Lub M, *et al.* 1995. *Immunol. Today* 16:479.
- Parsons J. 1996. *Curr. Opin. Cell Biol.* 8:146.

Related Products:

Product	Clone	Application
Purified anti-human CD11c	3.9	FC, IHC, CyTOF®
Purified anti-human CD11b (activated)	CBRM1/5	FC, IP
Purified anti-human CD11b	ICRF44	FC, IF, IHC, CyTOF®
Purified anti-human CD18	TS1/18	FC
Purified Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC, ICC, IF, IHC, IP, WB
APC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Biotin Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC, ELISA, IHC, IF, WB
FITC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
HRP Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	ELISA, IHC, WB
PE Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
Purified anti-human CD50	MEM-171	FC, IP



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.