

## **Product Data Sheet**

## FITC anti-human CD11b (activated)

Catalog # / Size: 301403 / 25 tests

301404 / 100 tests

Clone: CBRM1/5 **Isotype:** Mouse IgG1,  $\kappa$ 

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with

FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.


## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. Test size products are transitioning from 20 µl to 5 µl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.



PMA activated human granulocytes stained with CBRM1/5 FITC

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, and blocking<sup>1</sup> of the binding of ICAM-1 and fibrinogen. The CBRM1/5 antibody binds specifically to the activated form of CD11b. The LEAF™ purified

antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No.

301408).

- Application References: 1. Diamond M, et al. 1993. J. Cell Biol. 120:545. (Block)
  - 2. Bryn T, et al. 2006. J. Immunol. 176:7361.J. Immunol.
  - Sithu SD, et al. 2007. J. Biol. Chem. doi:10.1074/jbc.M611273200. 4. Barthel SR, et al. 2006. Am J. Respir Cell Mol Biol. 35:378. PubMed
  - 5. Pillay J,et al. 2010. J. Leukocyte Biol. 88:211. PubMed 6. Wilson RP, et al. 2011. Infect Immun. 79:830. PubMed 7. Reis ES, et al. 2012. J. Immunol. 189:4797. PubMed.

Description: The CBRM1/5 antibody reacts with an activated form of human CD11b, a 165-170 kD type I transmembrane glycoprotein also known as  $\alpha_M$  integrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin  $\beta_2$  (CD18) and is expressed on granulocytes, monocytes/macrophages dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils, as well as adhesion, phagocytosis and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen and Factor X. The CBRM1/5 antibody recognizes a subset of CD11b molecules on neutrophils and monocytes activated with chemoattractants or phorbol-esters. This antibody does not recognize "non-activated" CD11b. The epitope recognized by CBRM1/5 is contained in the I domain of the lpha chain. The CBRM1/5 antibody blocks cell adhesion to fibrinogen and ICAM-1.

- Antigen References: 1. Stewart M, et al. 1995. Curr. Opin. Cell Biol. 7:690.
  - 2. Petty H, et al. 1996. Immunol. Today 17:209.

Related Products: Product

Clone Application FITC anti-human CD11c 3.9 FC FC FITC anti-human CD11a HI111 PE anti-human CD11b ICRF44 TS1/18 FČ FITC anti-human CD18 FC, ICFC FC, ICC, ICFC FC, ICFC FITC Mouse IgG1, κ Isotype Ctrl MOPC-21 Cell Staining Buffer RBC Lysis Buffer (10X) Human TruStain FcX™ (Fc Receptor Blocking Solution) FC. ICC. ICFC



