

Product Data Sheet

APC anti-human CD11b (activated)

Catalog # / Size: 301409 / 25 tests

301410 / 100 tests

Clone: CBRM1/5 **Isotype:** Mouse IgG1, κ

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with

APC under optimal conditions. The solution is free of unconjugated APC and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



PMA-stimulated (5 minutes) human peripheral blood granulocytes stained with CBRM1/5 APC.

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. Test size products are transitioning from 20 μ l to 5 μ l per test. Please check your vial or your CoA to find the suggested use of this readth to per million cells in 100 μ l staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for

optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Additional reported applications (for the relevant formats) include: immunoprecipitation, and blocking¹ of the binding of Application Notes: ICAM-1 and fibrinogen. The CBRM1/5 antibody binds specifically to the activated form of CD11b. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No.

Application References: 1. Diamond M, et al. 1993. J. Cell Biol. 120:545. (Block)

 Bryn T, et al. 2006. J. Immunol. 176:7361.PubMed.
Sithu SD, et al. 2007. J. Biol. Chem. doi:10.1074/jbc.M611273200. Barthel SR, et al. 2006. Am J. Respir Cell Mol Biol. 35:378. PubMed

5. Pillay J, et al. 2010. J. Leukocyte Biol. 88:211. PubMed 6. Wilson RP, et al. 2011. Infect Immun. 79:830. PubMed

Description: The CBRM1/5 antibody reacts with an activated form of human CD11b, a 165-170 kD type I transmembrane glycoprotein also known as α_M integrin, Mac-1, CR3, and C3biR. CD11b non-covalently associates with integrin β_2 (CD18) and is expressed on granulocytes, monocytes/macrophages dendritic cells, NK cells, and subsets of T and B cells. CD11b/CD18 is critical for the transendothelial migration of monocytes and neutrophils, as well as adhesion, phagocytosis and neutrophil activation. CD11b/CD18 interacts with ICAM-1 (CD54), ICAM-2 (CD102), ICAM-4, CD14, CD23, heparin, iC3b, fibrinogen and Factor X. The CBRM1/5 antibody recognizes a subset of CD11b molecules on neutrophils and monocytes activated with chemoattractants or phorbol-esters. This antibody does not recognize "non-activated" CD11b. The epitope recognized by CBRM1/5 is contained in the I domain of the lpha chain. The CBRM1/5 antibody blocks cell adhesion to fibrinogen and ICAM-1.

Antigen References: 1. Stewart M, et al. 1995. Curr. Opin. Cell Biol. 7:690.

2. Petty H, et al. 1996. Immunol. Today 17:209.

Related Products: Product

APC Mouse IgG1, κ Isotype Ctrl

Cell Staining Buffer RBC Lysis Buffer (10X)

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone MOPC-21

Application FC, ICFC FC, ICC, ICFC FC, ICFC FC, ICC, ICFC



