

Biotin anti-human CD16

Catalog # / Size: 302003 / 25 µg
302004 / 100 µg

Clone: 3G8

Isotype: Mouse IgG1, κ

Workshop Number: V NK80


Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Pigtailed Macaque, Capuchin Monkey, Squirrel Monkey, Sooty Mangabey, Cotton-topped Tamarin, Marmoset

Preparation: The antibody was purified by affinity chromatography, and conjugated with biotin under optimal conditions. The solution is free of unconjugated biotin.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C. **Do not freeze.**



Human peripheral blood lymphocytes stained with biotinylated 3G8 and Sav-PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤ 1.0 µg per 10⁶ cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: The 3G8 antibody blocks neutrophil phagocytosis and stimulates NK cell proliferation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁶, immunoprecipitation³, stimulation of NK cell proliferation⁴, blocking of phagocytosis⁵, and blocking of immunoglobulin binding to FcγRIII^{7,8}. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302014). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302050) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Application References:

- Knapp W, *et al.* Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Edberg J, *et al.* 1997. *J. Immunol.* 159:3849. (IP)
- Hoshino S, *et al.* 1991. *Blood* 78:3232. (Stim)
- Tamm A, *et al.* 1996. *Immunol.* 157:1576. (Block)
- Da Silva DM, *et al.* 2001. *Int. Immunol.* 13:633. (IHC)
- Holl V, *et al.* 2004. *J. Immunol.* 173:6274. (Block)
- Hober D, *et al.* 2002. *J. Gen. Virol.* 83:2169. (Block)
- Brainard DM, *et al.* 2009. *J. Virol.* 83:7305. PubMed
- Smed-Sörensen A, *et al.* 2008. *Blood* 111:5037. (Block) PubMed
- Timmerman KL, *et al.* 2008. *J. Leukoc. Biol.* 84:1271. (FC) PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
- Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
- Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
- Boltz A, *et al.* 2011. *J. Biol Chem.* 286:21896. PubMed

Description: CD16 is known as low affinity IgG receptor III (FcγRIII). It is expressed as two distinct forms (CD16a and CD16b). CD16a (FcγRIIIA) is a 50-65 kD polypeptide-anchored transmembrane protein. It is expressed on the surface of NK cells, activated monocytes, macrophages, and placental trophoblasts in humans. CD16b (FcγRIIIB) is a 48 kD glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

Antigen References:

- Fleit H, *et al.* 1982. *P. Natl. Acad. Sci. USA* 79:3275.
- Stroncek D, *et al.* 1991. *Blood* 77:1572.
- Wirthmueller U, *et al.* 1992. *J. Exp. Med.* 175:1381.

Related Products:Product

Biotin anti-human CD64

PE anti-human CD32

Biotin Mouse IgG1, κ Isotype Ctrl

APC Streptavidin

APC/Cy7 Streptavidin

Clone

10.1

FUN-2

MOPC-21

Application

FC

FC

FC, ICFC

FC, ICFC

FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

FITC Streptavidin
HRP Streptavidin
PE Streptavidin
PE/Cy5 Streptavidin
PE/Cy7 Streptavidin
Cell Staining Buffer
RBC Lysis Buffer (10X)
PE anti-human CD56 (NCAM)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

MEM-188

FC, ICFC
ELISA, ELISPOT, IHC, WB
FC, ICFC
FC, ICFC
FC, ICFC
FC, ICC, ICFC
FC, ICFC
FC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.