

APC anti-human CD16

Catalog # / Size: 302011 / 25 tests
302012 / 100 tests

Clone: 3G8

Isotype: Mouse IgG1, κ


Workshop Number: V NK80

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Pigtailed Macaque, Capuchin Monkey, Squirrel Monkey, Sooty Mangabey, Cotton-topped Tamarin, Marmoset

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The CD16 antibody solution should be stored undiluted at 4°C, and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with 3G8 APC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The 3G8 antibody blocks neutrophil phagocytosis and stimulates NK cell proliferation. Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections⁶, immunoprecipitation³, stimulation of NK cell proliferation⁴, blocking of phagocytosis⁵, and blocking of immunoglobulin binding to Fc γ RIII^{7,8}. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 302014). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302050) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/ μ g).

- Application References:**
- Knapp W, *et al.* Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.
 - Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
 - Edberg J, *et al.* 1997. *J. Immunol.* 159:3849. (IP)
 - Hoshino S, *et al.* 1991. *Blood* 78:3232. (Stim)
 - Tamm A, *et al.* 1996. *Immunol.* 157:1576. (Block)
 - Da Silva DM, *et al.* 2001. *Int. Immunol.* 13:633. (IHC)
 - Holl V, *et al.* 2004. *J. Immunol.* 173:6274. (Block)
 - Hober D, *et al.* 2002. *J. Gen. Virol.* 83:2169. (Block)
 - Brainard DM, *et al.* 2009. *J. Virol.* 83:7305. PubMed
 - Smed-Sørensen A, *et al.* 2008. *Blood* 111:5037. (Block) PubMed
 - Timmerman KL, *et al.* 2008. *J. Leukoc. Biol.* 84:1271. (FC) PubMed
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
 - Kim WK, *et al.* 2006. *Am. J. Pathol.* 168:822. (FC)
 - Boltz A, *et al.* 2011. *J. Biol Chem.* 286:21896. PubMed

Description: CD16 is known as low affinity IgG receptor III (Fc γ RIII). It is expressed as two distinct forms (CD16a and CD16b). CD16a (Fc γ RIIIA) is a 50-65 kD polypeptide-anchored transmembrane protein. It is expressed on the surface of NK cells, activated monocytes, macrophages, and placental trophoblasts in humans. CD16b (Fc γ RIIIB) is a 48 kD glycosylphosphatidylinositol (GPI)-anchored protein. Its extracellular domain is over 95% homologous to that of CD16a, and it is expressed specifically on neutrophils. CD16 binds aggregated IgG or IgG-antigen complex which functions in NK cell activation, phagocytosis, and antibody-dependent cell-mediated cytotoxicity (ADCC).

- Antigen References:**
- Fleit H, *et al.* 1982. *P. Natl. Acad. Sci. USA* 79:3275.
 - Stroncek D, *et al.* 1991. *Blood* 77:1572.
 - Wirthmueller U, *et al.* 1992. *J. Exp. Med.* 175:1381.

Related Products: Product
PE anti-human CD64

Clone
10.1

Application
FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

APC anti-human CD32	FUN-2	FC
APC Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
APC anti-human CD56 (NCAM)	MEM-188	FC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.