

Purified anti-human CD18

Catalog # / Size: 302101 / 25 µg
302102 / 100 µg

Clone: TS1/18

Isotype: Mouse IgG1, κ

Workshop Number: V AS162


Reactivity: Human, **Cross-Reactivity:** Swine (Pig, Porcine)

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified TS1/18, followed by anti-mouse IgGs FITC

Applications:

Applications: FC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤0.25 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cell adhesion and migration^{3,4}. The LEAF™ Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302112). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302116) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Application References:

- Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York.
- Kishimoto T, *et al.* 1997. Leucocyte Typing VI. Garland Press. London.
- Van Epps DE, *et al.* 1989. *J. Immunol.* 143:3207. (Block)
- Meerschaert J, *et al.* 1994. *J. Immunol.* 152:1915. (Block)
- Sithu SD, *et al.* 2007. *J. Biol. Chem.* doi:10.1074/jbc.M611273200.
- Sommaggio R, *et al.* 2012. *J. Immunol.* 188:2075. PubMed.

Description: CD18 is a 90-95 kD type I transmembrane protein also known as integrin β₂ subunit, LFA-1 β subunit, and β₂ integrin. CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and associated α chains function in adhesion and signaling in hematopoietic cells.

Antigen References:

- Anderson D, *et al.* 1987. *Annu. Rev. Med.* 38:175.
- Springer T. 1994. *Cell* 76:301.

Related Products: Product

Product	Clone	Application
Purified anti-human CD11c	3.9	FC, IHC, CyTOF®
Purified anti-human CD11b (activated)	CBRM1/5	FC, IP
Purified anti-human CD11a	HI111	FC, IHC, WB
Purified anti-human CD11b	ICRF44	FC, IF, IHC, CyTOF®
Purified Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC, ICC, IF, IHC, IP, WB
APC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Biotin Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC, ELISA, IHC, IF, WB
PE Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.