

Product Data Sheet

PE/Cy5 anti-human CD18

Catalog # / Size: 302109 / 25 tests

302110 / 100 tests

Clone: TS1/18

Isotype: Mouse IgG1, κ

Workshop Number: V AS162

Reactivity: Human, Cross-Reactivity: Swine (Pig. Porcine)

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE/Cy5 under optimal conditions. The solution is free of unconjugated


PE/Cy5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes stained with TS1/18 PE/Cy5

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test

size products are transitioning from 20 µl to 5 µl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at

www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: inhibition of cell adhesion and migration^{3,4}. The

LEAF™ Purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 302112). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302116) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed

for research use only.

 Schlossman S, et al. 1995. Leucocyte Typing V. Oxford University Press. New York.
Kishimoto T, et al. 1997. Leucocyte Typing VI. Garland Press. London.
Van Epps DE, et al. 1989. J. Immunol. 143:3207. (Block) Application References: 1.

4. Meerschaert J, et al. 1994. J. Immunol. 152:1915. (Block) 5. Sithu SD, et al. 2007. J. Biol. Chem. doi:10.1074/jbc.M611273200.

6. Sommaggio R, et al. 2012. J. Immunol. 188:2075. PubMed.

Description: CD18 is a 90-95 kD type I transmembrane protein also known as integrin β_2 subunit, LFA-1 β subunit, and β_2 integrin.

CD18 non-covalently associates with CD11a, CD11b or CD11c. CD18 is expressed on all leukocytes. CD18 and

associated α chains function in adhesion and signaling in hematopoietic cells.

Antigen References: 1. Anderson D, et al. 1987. Annu. Rev. Med. 38:175.

2. Springer T. 1994. Cell 76:301.

Related Products: Product Clone Application

PE anti-human CD11c 3.9 PE anti-human CD11b (activated) CBRM1/5 FC FC PE/Cy5 anti-human CD11a HI111 PE/Cy5 anti-human CD11b PE/Cy5 Mouse IgG1, κ Isotype Ctrl ICRF44 MOPC-21 FC, ICFC

Cell Staining Buffer RBC Lysis Buffer (10X)

FC, ICC, ICFC FC, ICFC Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



