

## **Product Data Sheet**

## PE anti-human CD15 (SSEA-1)

Catalog # / Size: 323005 / 25 tests

323006 / 100 tests

Clone: W6D3

**Isotype:** Mouse IgG1,  $\kappa$ 

Immunogen: WERI-RB-1 retinoblastoma cell line

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE under optimal conditions. The solution is free of unconjugated PE and


unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood granulocytes stained with W6D3 PE

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 μl to 5 μl per test. Please check your vial or your CoA to find the

suggested use of this reagent per million cells in 100 µl staining volume or per 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at

www.biolegend.com/testsize regarding the test size change.

Description: CD15 is 3-fucosyl-N-acetyllactosamine (3-FAL) also known as Lewis X, 3-FAL, X-hapten, and SSEA-1. CD15 is

expressed on granulocytes and monocytes. It has also been shown to be expressed on Langerhans cells and some malignant cells. CD15 has been implicated in adhesion as well as chemotaxis, phagocytosis, and bactericidal activity.

Antigen References: 1. Stocks SC, et al. 1990. Biochem. J. 268:275.

**Related Products: Product** Clone Application

Cell Staining Buffer FC, ICC, ICFC FC MOPC-21

PE Mouse IgG1, κ Isotype Ctrl (FC) Human TruStain FcX™ (Fc Receptor Blocking Solution)



