

## **Product Data Sheet**

## Alexa Fluor® 647 anti-human CD19

Catalog # / Size: 302222 / 25 tests

302220 / 100 tests

Clone: HIB19

**Isotype:** Mouse IgG1,  $\kappa$ 

Workshop Number: V CD19.11

Reactivity: Human, Cross-Reactivity: Chimpanzee

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 647 under optimal conditions. The solution is free of


unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes stained with HIB19 Alexa Flour® 647

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 µl per million cells in 100 µl volume or 5 µl per 100 ul of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

\* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at 633nm / 635nm.

\*\* Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application Notes: Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections and blocking of B cell proliferation. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is recommended for functional assays (Cat. No. 302214).

- Application References: 1. Schlossman S, et al. 1995. Leucocyte Typing V. Oxford University Press. New York. 2. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press. New York.
  - 3. Bradbury L, et al. 1993. J. Immunol. 151:2915. 4. Joseph A, et al. 2010. J. Virol. 84:6645. PubMed
  - 5. Wang X, et al. 2010. Haematologica. 95:884. (FC) PubMed 6. Walker JD, et al. 2009. J. Immunol. 182:1548. (Block) PubMed
  - 7. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC) 8. Geneve L, et al. 2012. Int Immunol. 24:645. PubMed.

**Description:** CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cells, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.

Antigen References: 1. Tedder T, et al. 1994. Immunol. Today 15:437.

2. Bradbury L, et al. 1993. J. Immunol. 151:2915.

**Related Products: Product** Clone Application FC, ICC, ICFC

Cell Staining Buffer

RBC Lysis Buffer (10X) Alexa Fluor® 647 Mouse IgG1, κ Isotype Ctrl (FC) FC, ICFC FC, IF MOPC-21 Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



