

## **Product Data Sheet**

## Alexa Fluor® 700 anti-human CD19

Catalog # / Size: 302225 / 25 µg

302226 / 100 µg

Clone: HIB19

**Isotype:** Mouse IgG1,  $\kappa$ 

Workshop Number: V CD19.11

Reactivity: Human, Cross-Reactivity: Chimpanzee

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 700 under optimal conditions. The solution is free of


unconjugated Alexa Fluor® 700.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes stained with HIB19 Alexa Fluor® 700

## **Applications:**

Applications: FC - Quality tested

Recommended Usage: This reagent is developed for immunofluorescent staining for flow cytometric analysis; the suggested use of this reagent is ≤1.0 µg per million cells in 100 µl volume. It is highly recommended that the reagent be titrated for optimal performance for each application.

> \* Alexa Fluor® 700 has a maximum emission of 719 nm when it is excited at 633 nm / 635 nm. Prior to using Alexa Fluor® 700 conjugate for flow cytometric analysis, please verify your flow cytometer's capability of exciting and detecting the fluorochrome.

> Alexa Fluor® 700 is a registered trademark of Molecular Probes, Inc. Alexa Fluor® 700 dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with

microarrays and high content screening, and are covered by pending and issued patents.

**Application Notes:** 

Additional reported applications (for the relevant formats) include: immunohistochemical staining of acetone-fixed frozen tissue sections and blocking of B cell proliferation. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302214).

- **Application References:** 1. Schlossman S, *et al.* 1995. Leucocyte Typing V. Oxford University Press. New York. 2. Knapp W, *et al.* 1989. Leucocyte Typing IV. Oxford University Press. New York.
  - 3. Bradbury L, et al. 1993. J. Immunol. 151:2915.
  - 4. Joseph A, et al. 2010. J. Virol. 84:6645. PubMed 5. Wang X, et al. 2010. Haematologica. 95:884. (FC) PubMed 6. Walker JD, et al. 2009. J. Immunol. 182:1548. (Block) PubMed
  - 7. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)

Description: CD19 is a 95 kD type I transmembrane glycoprotein also known as B4. It is a member of the immunoglobulin superfamily expressed on B-cells (from pro-B to blastoid B cells, absent on plasma cells) and follicular dendritic cells. CD19 is involved in B cell development, activation, and differentiation. CD19 forms a complex with CD21 (CR2) and CD81 (TAPA-1), and functions as a BCR co-receptor.

Antigen References: 1. Tedder T, et al. 1994. Immunol. Today 15:437.

2. Bradbury L, et al. 1993. J. Immunol. 151:2915.

**Related Products: Product** Clone

**Application** FC, ICC, ICFC FC, ICFC Cell Staining Buffer RBC Lysis Buffer (10X) Alexa Fluor® 700 Mouse IgG1, κ Isotype Ctrl MOPC-21 FC, ICFC Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



