

Product Data Sheet

Alexa Fluor® 488 anti-human CD20

Catalog # / Size: 302316 / 100 tests

Clone: 2H7

Isotype: Mouse IgG2b, κ

Workshop Number: IV B201

Reactivity: Human, Cross-Reactivity: Chimpanzee, Baboon, Cynomolgus, Rhesus,

Pigtailed Macaque, Capuchin Monkey, Squirrel Monkey

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 488 under optimal conditions. The solution is free of


unconjugated Alexa Fluor® 488.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes were stained with anti-CD20 (clone 2H7) Alexa Fluor® 488 (filled histogram), or mouse IgG2b, κ Alexa Fluor® 488 (open histogram).

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μl per million cells or 5 μl per 100 μl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

- * Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at 488 nm.
- ** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Fluor® dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and issued patents.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation⁴ and immunohistochemical staining of acetone-fixed frozen sections⁵.

- Application References: 1. Schlossman S, et al. 1995. Leucocyte Typing V. Oxford University Press. New York. 2. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press. New York. 3. McMichael A, et al. 2004. Blood 99:3256. (IP)

 - 5. Mack CL, et al. 2004. Pediatr. Res. 56:79. (IHC)

Description: CD20 is a 33-37 kD, four transmembrane spanning protein, also known as B1 and Bp35. CD20 is expressed on pre-B-cells, resting and activated B cells (not plasma cells), some follicular dendritic cells, and at low levels on a T cell subset. CD20 is heavily phosphorylated on activated B cells and malignant B cells. Homo-oligomeric complexes of CD20 are thought to form Ca2+ conductive ion channels in the plasma membrane of B cells. The CD20 molecule is involved in B-cell activation and is associated with various Src family kinases (Lyn, Lck, Fyn). It exists in a complex with MHC class I and II, CD53, CD81, and CD82.

Antigen References: 1. Hultin L, et al. 1993. Cytometry 14:196.

2. Tedder T, et al. 1994. Immunol. Today 15:450.

Related Products: Product

Cell Staining Buffer Alexa Fluor® 488 Mouse IgG2b, κ Isotype Ctrl Human TruStain FcX™ (Fc Receptor Blocking Solution) Clone MPC-11

FC, ICC, ICFC FC, ICFC

Application



