


Anti-Mouse CD14 PerCP-Cyanine5.5

Catalog Number: 45-0141

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of mouse thioglycolate-elicited peritoneal exudate cells with 0.25 ug of Rat IgG2a K Isotype Control PerCP-Cyanine5.5 (cat. 45-4321) (blue histogram) or 0.25 ug of Anti-Mouse CD14 PerCP-Cyanine5.5 (purple histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD14 PerCP-Cyanine5.5

REF Catalog Number: 45-0141

Clone: Sa2-8

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, kappa Formulation: aqueous buffer, 0.09% sodium azide, may contain

carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light

sensitive material.

Eatch Code: Refer to Vial

Use By: Refer to Vial

Contains sodium azide

Description

The Sa2-8 monoclonal antibody reacts with mouse CD14, a 53-55 kDa GPI-linked glycoprotein. CD14 is a receptor for the complexes of LPS and LBP (LPS-Binding Protein) and is shown to associate with ToII-Like Receptor 4 (TLR4) and participate in the signaling and cellular response to bacterial LPS. In mouse, CD14 is expressed on the surface of macrophages and under certain conditions is also found in the serum in a secreted form. Sa2-8 has weak antagonistic activity (in NF-kappaB activation or TNF alpha production with LPS stimulation).

Applications Reported

This Sa2-8 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This Sa2-8 antibody has been tested by flow cytometric analysis of mouse thioglycolate-elicited peritoneal exudate cells. This can be used at less than or equal to 0.5 μ g per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Akashi S, Saitoh S, Wakabayashi Y, Kikuchi T, Takamura N, Nagai Y, Kusumoto Y, Fukase K, Kusumoto S, Adachi Y, Kosugi A, Miyake K. Lipopolysaccharide interaction with cell surface Toll-like receptor 4-MD-2: higher affinity than that with MD-2 or CD14. J Exp Med. 2003 198(7): 1035-42.

Related Products

45-0149 Anti-Human CD14 PerCP-Cyanine5.5 (61D3) 45-4321 Rat IgG2a K Isotype Control PerCP-Cyanine5.5 (eBR2a)

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT #5,268,486, 5,59,597 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.