


An Affymetrix Company

Anti-Mouse CD25 PerCP-Cyanine5.5

Catalog Number: 45-0251

Also known as: Interleukin-2 Receptor alpha

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of SJL splenocytes with Anti-Mouse CD4 FITC (cat. 11-0042) and 0.125 ug of Rat IgG1 K Isotype Control PerCP-Cyanine 5.5 (cat. 45-4301) (left) or 0.125 ug of Anti-Mouse CD25 PerCP-Cyanine 5.5 (right). Cells in the lymphocyte gate were used for analysis.

Product Information

Contents: Anti-Mouse CD25 PerCP-

Cyanine5.5

REF Catalog Number: 45-0251


Clone: PC61.5

Concentration: 0.2 mg/mL Host/Isotype: Rat IgG1, lambda



Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer Temperature Limitation: Store at 2-8°C. Do not

freeze. Light sensitive material. Batch Code: Refer to vial





Description

The PC61.5 antibody reacts with mouse CD25, the 55 kDa interleukin-2 receptor alpha chain (IL-2R alpha). CD25 is expressed by early progenitors of the T and B lineage as well as by activated mature T and B lymphocytes. By itself, CD25 binds IL-2 only with low affinity. However, CD25 associates with CD122 (IL-2 receptor beta chain) and CD132 (common gamma chain) to form the high affinity IL-2 receptor. Binding of IL-2 to both the high and low affinity classes of IL-2 receptor is inhibited by the PC61.5 antibody. CD25 plays a role in lymphocyte differentiation and activation/proliferation.

Applications Reported

This PC61.5 antibody has been reported for use in flow cytometric analysis.

Applications Tested

This PC61.5 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.25 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Tuve S, Chen BM, Liu Y, Cheng TL, Touré P, Sow PS, Feng Q, Kiviat N, Strauss R, Ni S, Li ZY, Roffler SR, Lieber A. Combination of tumor site-located CTL-associated antigen-4 blockade and systemic regulatory T-cell depletion induces tumor-destructive immune responses. Cancer Res. 2007 Jun 15;67(12):5929-39. (PC61.5, in vivo depletion, PubMed)

Huang B, Zhao J, Shen S, Li H, He KL, Shen GX, Mayer L, Unkeless J, Li D, Yuan Y, Zhang GM, Xiong H, Feng ZH. Listeria monocytogenes promotes tumor growth via tumor cell toll-like receptor 2 signaling. Cancer Res. 2007 May 1;67(9):4346-52. (**PC61.5**, in vivo depletion, PubMed)

Hayashi T, Hasegawa K, Adachi C. Elimination of CD4+CD25+ T cell accelerates the development of



An Affymetrix Company

Anti-Mouse CD25 PerCP-Cyanine5.5

Catalog Number: 45-0251

Also known as: Interleukin-2 Receptor alpha

RUO: For Research Use Only. Not for use in diagnostic procedures.

glomerulonephritis during the preactive phase in autoimmune-prone female NZB × NZW F1 mice. Int J Exp Pathol. 2005 Oct;86(5):289-96. (**PC61.5**, in vivo depletion, PubMed)

Lowenthal JW, Corthésy P, Tougne C, Lees R, MacDonald HR, Nabholz M. High and low affinity IL 2 receptors: analysis by IL 2 dissociation rate and reactivity with monoclonal anti-receptor antibody PC61. J Immunol. 1985 Dec;135(6):3988-94.

Lowenthal JW, Zubler RH, Nabholz M, MacDonald HR. Similarities between interleukin-2 receptor number and affinity on activated B and T lymphocytes. Nature. 1985 Jun 20-26;315(6021):669-72.

Lowenthal JW, Tougne C, MacDonald HR, Smith KA, Nabholz M. Antigenic stimulation regulates the expression of IL 2 receptors in a cytolytic T lymphocyte clone. J Immunol. 1985 Feb;134(2):931-9.

Related Products

00-4300 10X RBC Lysis Buffer (Multi-species) 11-0042 Anti-Mouse CD4 FITC (RM4-5) 12-5773 Anti-Mouse/Rat Foxp3 PE (FJK-16s) 45-4301 Rat IgG1 K Isotype Control PerCP-Cyanine5.5

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486, 5,569,587 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 2. suce of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.