


# **Anti-Mouse CD69 PerCP-Cyanine5.5**

Catalog Number: 45-0691

Also Known As: Very Early Activation Antigen, VEA

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of 1-day ConA-stimulated BALB/c splenocytes with 0.25 ug of Armenian Hamster IgG Isotype Control PerCP-Cyanine5.5 (cat. 45-4888) (open histogram) or 0.25 ug of Anti-Mouse CD69 PerCP-Cyanine5.5 (filled histogram). Total viable cells were used for analysis.

# **Product Information**

Contents: Anti-Mouse CD69 PerCP-Cyanine5.5

REF Catalog Number: 45-0691

Clone: H1.2F3

Concentration: 0.2 mg/mL

Host/Isotype: Armenian Hamster IgG

Formulation: aqueous buffer, 0.09% sodium azide, may contain

carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light

sensitive material.

Batch Code: Refer to Vial

Use By: Refer to Vial

Caution, contains Azide

## Description

The H1.2F3 monoclonal antibody reacts with mouse CD69, also known as very early activation antigen (VEA). CD69 is approximately 35 kDa and is expressed on the surface as a disulfide-linked dimer. While a small subset of lymphocytes in the thymus, spleen and lymph nodes express this antigen, activation of both T and B cells rapidly upregulates the surface expression of CD69, suggesting a role for CD69 in lymphocyte development and activation.

#### **Applications Reported**

This H1.2F3 antibody has been reported for use in flow cytometric analysis.

### **Applications Tested**

This H1.2F3 antibody has been tested by flow cytometric analysis of activated splenocytes. This can be used at less than or equal to 0.5  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

# References

Yokoyama, W. M., F. Koning, et al. (1988). Characterization of a cell surface-expressed disulfide-linked dimer involved in murine T cell activation. J Immunol 141(2): 369-76.

#### **Related Products**

45-4888 Armenian Hamster IgG Isotype Control PerCP-Cyanine5.5 (eBio299Arm)

#### Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486, 5,569,587 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 3. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.