


Anti-Human Foxp3 PerCP-Cyanine5.5

Catalog Number: 45-4776 Also known as: Forkhead Box P3, Scurfin, JM2 RUO: For Research Use Only. Not for use in diagnostic procedures.



X

LOT

REF Catalog Number: 45-4776 Clone: PCH101 Concentration: 5 uL (0.125 ug)/test Host/Isotype: Rat IgG2a, kappa Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material. Batch Code: Refer to vial Use By: Refer to vial Contains sodium azide

Description

eBioscience offers a panel of monoclonal antibodies to different epitopes of human Foxp3, providing useful tools for investigating the complete expression pattern of Foxp3 at the protein level, and discerning the precise subsets of Foxp3⁺ cells. Please contact tech@ebioscience.com or 888.810.6168 for any additional assistance.

The PCH101 antibody reacts with the amino terminus of human foxp3 protein also known as FORKHEAD BOX P3, SCURFIN, and JM2; cross reactivity of this antibody to other proteins has not been determined. Foxp3, a 49-55 kDa protein, is a member of the forkhead/winged-helix family of transcriptional regulators, and was identified as the gene defective in 'scurfy' (sf) mice. Constitutive high expression of Foxp3 mRNA has been shown in CD4+CD25+ regulatory T cells (Treg cells), and ectopic expression of foxp3 in CD4+CD25- cells imparts a Treg phenotype in these cells.

Intracellular staining of human peripheral blood mononuclear cells (PBMCs) with PCH101 antibody using the antihuman Foxp3 Staining Set and protocol reveals approximately 0.5-4% of lymphocytes staining, with the majority of staining occurring in the CD25^{bright} population. This is subject to donor variability.

PCH101 crossreacts with rhesus, chimpanzee and cynomolgus. We recommend the use of CD4 (OKT4, cat. 11-0048, or RPA-T4, cat. 11-0049, depending on the species) and CD25 (BC96, cat. 17-0259).

Applications Reported

This PCH101 antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

Applications Tested

This PCH101 antibody has been pre-titrated and tested by intracellular staining and flow cytometric analysis of normal human peripheral blood cells using the Foxp3 Buffers and protocol. Please refer to Best Protocols for Staining Protocol (refer to Protocol B: One-step protocol for intracellular (nuclear) proteins). This can be used at 5 μ L (0.125 μ g) per test. A test is defined as the amount (μ g) of antibody that will stain a cell sample in a final volume of 100 μ L. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test.



Anti-Human Foxp3 PerCP-Cyanine5.5

Catalog Number: 45-4776 Also known as: Forkhead Box P3, Scurfin, JM2 RUO: For Research Use Only. Not for use in diagnostic procedures.

References

Bell MP, Svingen PA, Rahman MK, Xiong Y, Faubion WA Jr. FOXP3 regulates TLR10 expression in human T regulatory cells. J Immunol. 2007 Aug 1;179(3):1893-900. (**PCH101**, ChIP assay, PubMed)

Clark RA, Kupper TS. IL-15 and dermal fibroblasts induce proliferation of natural regulatory cells isolated from human skin. Blood. 2007 Jan 1;109(1):194-202. (**PCH101**, IHC frozen, PubMed)

Epple HJ, Loddenkemper C, Kunkel D, Tröger H, Maul J, Moos V, Berg E, Ullrich R, Schulzke JD, Stein H, Duchmann R, Zeitz M, Schneider T. Mucosal but not peripheral FOXP3+ regulatory T cells are highly increased in untreated HIV infection and normalize after suppressive HAART. Blood. 2006 Nov 1;108(9):3072-8. **PCH101**, IHC paraffin, PubMed)

Manigold T, Shin EC, Mizukoshi E, Mihalik K, Murthy KK, Rice CM, Piccirillo CA, Rehermann B. Foxp3+CD4+CD25+ T cells control virus-specific memory T cells in chimpanzees recovered from Hepatitis C.Blood. 2006 Jun 1;107(11):4424-32. (**PCH101** in chimpanzee, PubMed)

Ahmadzadeh M, Rosenberg SA. IL-2 Administration Increases CD4+CD25hiFoxp3+ Regulatory T Cells in Cancer Patients. Blood. 2006 Mar 15;107(6):2409-14. (PCH101, IC Flow, PubMed)

Hartwig UF, Nonn M, Khan S, Meyer RG, Huber C, Herr W. Depletion of alloreactive T cells via CD69: implications on antiviral, antileukemic and immunoregulatory T lymphocytes. Bone Marrow Transplant. 2006 Feb;37(3):297-305. (PCH101, IC Flow, PubMed)

Crellin NK, Garcia RV, Hadisfar O, Allan SE, Steiner TS, Levings MK. Human CD4+ T Cells Express TLR5 and Its Ligand Flagellin Enhances the Suppressive Capacity and Expression of FOXP3 in CD4+CD25+ T Regulatory Cells. J Immunol. 2005 Dec 15;175(12):8051-9. (**PCH101**, IC Flow, PubMed)

Lim, H.W., P. Hillsamer, A.H. Banham, and C.H. Kim. 2005. Cutting Edge: Direct Suppression of B cells by CD4+CD25+ Regulatory T cells. J Immunol. 2005 Oct 1;175(7):4180-3. (**PCH101**,IC Flow, PubMed)

Manavalan JS, Kim-Schulze S, Scotto L, Naiyer AJ, Vlad G, Colombo PC, Marboe C, Mancini D, Cortesini R, Suciu-Foca N. Alloantigen specific CD8+CD28- FOXP3+ T suppressor cells induce ILT3+ ILT4+ tolerogenic endothelial cells, inhibiting alloreactivity. Int Immunol. 2004 Aug;16(8):1055-68.

Takahata Y, Nomura A, Takada H, Ohga S, Furuno K, Hikino S, Nakayama H, Sakaguchi S, Hara T. CD25+CD4+ T cells in human cord blood: an immunoregulatory subset with naïve phenotype and specific expression of forkhead box p3 (Foxp3) gene. Exp Hematol. 2004 Jul;32(7):622-9.

Sakaguchi S. 2003. The origin of FOXP3-expressing CD4+ regulatory T cells: thymus or periphery. J Clin Invest. 2003 November 1; 112(9): 1310–1312.

Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. Science. 2003 Feb 14;299(5609):1057-61.

Related Products

00-5523 Foxp3 / Transcription Factor Staining Buffer Set 12-0259 Anti-Human CD25 PE (BC96) 17-0048 Anti-Human CD4 APC (OKT4 (OKT-4)) 45-4321 Rat IgG2a K Isotype Control PerCP-Cyanine5.5 (eBR2a) 45-5773 Anti-Mouse/Rat Foxp3 PerCP-Cyanine5.5 (FJK-16s)



Anti-Human Foxp3 PerCP-Cyanine5.5

Catalog Number: 45-4776 Also known as: Forkhead Box P3, Scurfin, JM2 RUO: For Research Use Only. Not for use in diagnostic procedures.

Legal

FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT FOR NON-COMMERCIAL RESEARCH USE ONLY. NOT FOR THERAPEUTIC OR IN VIVO APPLICATIONS. OTHER USE NEEDS LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. UNDER U.S. PATENT # 5,268,486,559,597 AND 5,627,027 AND FOREIGN EQUIVALENTS AND PENDING APPLICATIONS. THIS MATERIAL IS SUBJECT TO PROPRIETARY RIGHTS OF GE HEALTHCARE BIO-SCIENCES CORP. AND CARNEGIE MELLON UNIVERSITY AND MADE AND SOLD UNDER LICENSE FROM GE HEALTHCARE BIO-SCIENCES CORP. THIS PRODUCT IS LICENSED FOR SALE ONLY FOR RESEARCH. IT IS NOT LICENSED FOR ANY OTHER USE. THERE IS NO IMPLIED LICENSE HEREUNDER FOR ANY COMMERCIAL USE. COMMERCIAL USE shall include: 1. sale, lease, license or other transfer of the material or any material derived or produced from it; 2. sale, lease, license or other grant of rights to use this Material or any material derived or produced from it; 2. use of this material to perform services for a fee for third parties. IF YOU REQUIRE A COMMERCIAL LICENSE TO USE THIS MATERIAL, AND DO NOT HAVE ONE, RETURN THIS MATERIAL, UNOPENED TO EBIOSCIENCE, INC. 10255 SCIENCE CENTER DRIVE, SAN DIEGO, CALIFORNIA 92121 USA AND ANY MONEY PAID FOR THE MATERIAL WILL BE REFUNDED.