

## **Product Data Sheet**

104

Purified anti-human CD27

Catalog # / Size:	302802 / 100 µg						
Clone:	O323		i.				
Isotype:	Mouse IgG1, κ		11				
Workshop Number:	IV T-186	5	M		1		
Reactivity:	Human, Cross-Reactivity: Baboon, Cynomolgus, Rhesus, Squirrel Monkey	Number	. IN				
Preparation:	The antibody was purified by affinity chromatography.	Cell N	14				
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.		11				
Concentration:	0.5 mg/ml	Relative	, "N				
Storage:	The antibody solution should be stored undiluted at 4°C.		y -	6 6	M .		
Application	S:	10	ρ	10 <sup>1</sup>	10 <sup>2</sup>	10 <sup>3</sup>	10
Applications:	FC - Quality tested CyTOF® - Validated	Hı	uman ı	2	resence In al blood	itensity	cvtes
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent		stained with purified 0323, followe				

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq$ 1.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References: 1. Knapp W, et al. Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.

Description: CD27 is a 50-55 kD type I membrane protein also known as S152 and T14. It is a lymphocyte-specific member of the TNF-receptor superfamily. CD27 is expressed on medullary thymocytes, virtually all mature T cells, some B cells, and NK cells. CD27 binds to CD70 and plays an important role in costimulation of T cell activation, and regulation of B cell differentiation and proliferation. The cytoplasmic domains of CD27 have also been shown to interact with TRAF2 and TRAF5 to elicit NF-kB and SAPK/JNK activation.

Antigen References: 1. Hintzen R, et al. 1994. Immunol. Today 15:307. 2. Agematsu K, et al. 1995. J. Immunol. 154:3627.

## **Related Products: Product**

Purified Mouse IgG1,  $\kappa$  Isotype Ctrl Cell Staining Buffer RBC Lysis Buffer (10X)

Clone MOPC-21

Application FC, ICFC, ICC, IF, IHC, IP, WB FC, ICC, ICFC FC, ICFC

by anti-mouse IgGs FITC



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.