

APC anti-human CD28

Catalog # / Size: 302911 / 25 tests
302912 / 100 tests

Clone: CD28.2

Isotype: Mouse IgG1, κ


Workshop Number: V-CD28.05

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Pigtailed Macaque, Squirrel Monkey, Capuchin Monkey, Sooty Mangabey

Preparation: The antibody was purified by affinity chromatography, and conjugated with APC under optimal conditions. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with APC CD28.2

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, immunohistochemical staining of acetone-fixed frozen tissue sections⁴, and *in vitro* T cell costimulation⁵⁻⁸. The CD28.2 antibody co-stimulates T cell proliferation and cytokine production in the presence of suboptimal amounts of anti-CD3 antibody. The LEAF™ purified antibody (Endotoxin <0.1 EU/ μ g, Azide-Free, 0.2 μ m filtered) is recommended for functional assays (Cat. No. 302914). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302934) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/ μ g).

Application References:

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- Nunes J, *et al.* 1993. *Biochem. J.* 293:835.
- Calea-Lauri J, *et al.* 1999. *J. Immunol.* 163:62.
- Tazi A, *et al.* 1999. *J. Immunol.* 163:3511. (IHC)
- Marti F, *et al.* 2001. *J. Immunol.* 166:197. (Costim)
- Jeong SH, *et al.* 2004. *J. Virol.* 78:6995. (Costim)
- Rivollier A, *et al.* 2004. *Blood* 104:4029. (Costim)
- Scharschmidt E, *et al.* 2004. *Mol. Cell Biol.* 24:3860. (Costim)
- Sheng W, *et al.* 2007. *Elsevier* 580:6819. PubMed
- Mitsuhashi M. 2007. *Clin Chem.* 53:148. PubMed
- Ye Z, *et al.* 2008. *Infect. Immun.* 76:2541. PubMed
- Magatti M, *et al.* 2008. *Stem Cells* 26:182. (FA) PubMed
- Yoshino N, *et al.* 2000. *Exp. Anim.* (Tokyo) 49:97. (FC)
- Rout N, *et al.* 2010. *PLoS One* 5:e9787. (FC)
- Leonard JA, *et al.* 2011. *J. Virol.* 85:6867. PubMed

Description: CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. *In vitro* studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell activation and proliferation.

Antigen References:

- Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
- June CH, *et al.* 1994. *Immunol. Today* 15:321.
- Linskey PS, *et al.* 1993. *Annu. Rev. Immunol.* 11:191.

Related Products:

Product
PE anti-human CD80

PE anti-human CD86

APC Mouse IgG1, κ Isotype Ctrl

Clone

2D10

IT2.2

MOPC-21

Application

FC

FC

FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Cell Staining Buffer
RBC Lysis Buffer (10X)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 9727 Pacific Heights Blvd, San Diego, CA 92121 www.biolegend.com
Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587