(1)



Product Data Sheet

Alexa Fluor® 488 anti-human CD28

Catalog # / Size:	302915 / 25 tests 302916 / 100 tests	
Clone:	CD28.2	h.
Isotype:	Mouse IgG1, κ	N. I
Workshop Number:	V-CD28.05	ž 11 i
Reactivity:	Human, Cross-Reactivity: Chimpanzee, Baboon, Cynomolgus, Rhesus, Pigtailed Macaque, Squirrel Monkey, Capuchin Monkey, Sooty Mangabey	Relative Cell Number
Preparation:	The antibody was purified by affinity chromatography, and conjugated with Alexa Fluor® 488 under optimal conditions. The solution is free of unconjugated Alexa Fluor® 488.	Lead the
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	V. V. N.
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.	10 ⁰ 10 ¹ 10 ² 10 ³ 10 ⁴ Log Fluoresence Intensity
Application	IS:	Human peripheral blood lymphocytes stained with CD28.2 Alexa Fluor® 488
Applications:	FC - Quality tested	
Recommended Usage:	Each lot of this antibody is quality control tested by immunofluorescent staining immunofluorescent staining, the suggested use of this reagent is 5 μ l per mill blood. It is recommended that the reagent be titrated for optimal performance	ion cells or 5 µl per 100 µl of whole
	* Alexa Fluor® 488 has a maximum emission of 519 nm when it is excited at ** Alexa Fluor® is a registered trademark of Molecular Probes, Inc. Alexa Flu under license from Molecular Probes, Inc. for research use only, except for us high content screening, and are covered by pending and issued patents.	or® dye antibody conjugates are sold
Application Notes:	Additional reported applications (for the relevant formats) include: immunoprecipitation, immunohistochemical staining of acetone-fixed frozen tissue sections ⁴ , and <i>in vitro</i> T cell costimulation ⁵⁻⁸ . The CD28.2 antibody co-stimulates T cell proliferation and cytokine production in the presence of suboptimal amounts of anti-CD3 antibody. The LEAF™ purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 302914). For highly sensitive assays, we recommend Ultra-LEAF™ purified antibody (Cat. No. 302934) with a lower endotoxin limit than standard LEAF™ purified antibodies (Endotoxin <0.01 EU/µg).	
Application References:	 Schlossman S, <i>et al.</i> Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. Nunes J, <i>et al.</i> 1993. <i>Biochem. J.</i> 293:835. Calea-Lauri J, <i>et al.</i> 1999. <i>J. Immunol.</i> 163:62. Tazi A, <i>et al.</i> 1999. <i>J. Immunol.</i> 163:3511. (IHC) Marti F, <i>et al.</i> 2001. <i>J. Immunol.</i> 166:197. (Costim) Jeong SH, <i>et al.</i> 2004. <i>J. Virol.</i> 78:6995. (Costim) Scharschmidt E, <i>et al.</i> 2004. <i>Biood</i> 104:4029. (Costim) Scharschmidt E, <i>et al.</i> 2004. <i>Mol. Cell Biol.</i> 24:3860. (Costim) Sheng W, <i>et al.</i> 2007. <i>Elsevier</i> 580:6819. PubMed Mitsuhashi M. 2007. <i>Clin Chem.</i>53:148. PubMed Ye Z, <i>et al.</i> 2008. <i>Infect. Immun.</i> 76:2541. PubMed Yoshino N, <i>et al.</i> 2008. <i>Stem Cells</i> 26:182. (FA) PubMed Yoshino N, <i>et al.</i> 2010. Exp. Anim. (Tokyo) 49:97. (FC) Rout N, <i>et al.</i> 2011. <i>J. Virol.</i> 85:6867. PubMed 	
Description:	CD28 is a 44 kD disulfide-linked homodimeric type I glycoprotein. It is a member of the immunoglobulin superfamily and is also known as T44 or Tp44. CD28 is expressed on most T lineage cells, NK cell subsets, and plasma cells. CD28 binds both CD80 and CD86 using a highly conserved motif MYPPY in the CDR3-like loop. CD28 is considered a major co-stimulatory molecule, inducing T lymphocyte activation and IL-2 synthesis, and preventing cell death. <i>In</i> <i>vitro</i> studies indicate that ligation of CD28 on T cells by CD80 and CD86 on antigen presenting cells provides a costimulatory signal required for T cell activation and proliferation.	
Antinan Defenses	4. Cablessmen C. et al. Eds. 1005 Lawssevite Typing V/ Ovfard University Dr.	and New York

Antigen References: 1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. 2. June CH, *et al.* 1994. *Immunol. Today* 15:321. 3. Linskey PS, *et al.* 1993. *Annu. Rev. Immunol.* 11:191. Clone Application

Related Products: Product

 (\underline{n})

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

Cell Staining Buffer		FC, ICC, ICFC
Alexa Fluor® 488 Mouse IgG1, κ Isotype Ctrl (FC)	MOPC-21	FC, IF
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.

BioLegend Inc., 9727 Pacific Heights Blvd, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587