

Product Data Sheet

PE/Cy7 anti-human CD31

Catalog # / Size:	303117 / 25 tests 303118 / 100 tests	8	
Clone:	WM59	2	a a
Isotype:	Mouse IgG1, κ	8	
Workshop Number:	V P025	τ.	
Reactivity:	Human, Cross-Reactivity: African Green, Baboon, Cynomolgus, Rhesus	Scatte	
Preparation:	The antibody was purified by affinity chromatography, and conjugated with PE/Cy7 under optimal conditions. The solution is free of unconjugated PE/Cy7 and unconjugated antibody.	Side S	
Formulation:	Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).	8	20
Storage:	The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. Do not freeze.		WM59 PE/Cy7
		Hı	ıman peripheral blood lvn

Applications:

Human peripheral blood lymphocytes, monocytes and granulocytes were stained with CD31 (clone WM59) PE/Cy7.

Applications: FC - Quality tested Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test size products are transitioning from 20 µl to 5 µl per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change. Application Notes: Additional reported applications (for the relevant formats) include: immunofluorescence microscopy², immunohistochemical staining of acetone-fixed frozen tissue sections, and blocking of platelet aggregation³. Clone WM59 is not recommended for immunohistochemical staining of formalin-fixed paraffin-embedded sections. The LEAF™ Purified antibody (Endotoxin <0.1 EU/µg, Azide-Free, 0.2 µm filtered) is recommended for functional assays (Cat. No. 303108). Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed for research use only. Application References: 1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V Oxford University Press. New York. 2. Muczynski KA, *et al.* 2003. *J. Am. Soc. Nephrol.* 14:1336. (IF) Wu XW, et al. 1997. Arterioscl. Throm. Vas. 17:3154. (Block)
Nagano M, et al. 2007. Blood 110:151. (FC) PubMed
MacFadyen JR, et al. 2005. FEBS Lett. 579:2569. PubMed 6. Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC) 7. Sestak K, et al. 2007. Vet. Immunol. Immunopathol. 119:21. 8. Wicki A, *et al.* 2012. *Clin Cancer Res.* 18:454. PubMed 9. Oeztuerk-Winder F, *et al.* 2012. *EMBO J.* 31:3431. (FC) PubMed 10. Ardehali R, *et al.* 2013. *PNAS*. PubMed. Description: CD31 is a 130-140 kD type I transmembrane glycoprotein also known as platelet endothelial cell adhesion molecule-1 (PECAM-1) or Endocam. It is expressed on monocytes, platelets, granulocytes, endothelial cells and lymphocyte subsets. CD31 has been reported to bind CD38 and be involved in wound healing, angiogenesis, and cellular migration in an inflammatory situation. Clone WM59 has been reported to recognize the D2 extracellular portion of CD31. Antigen References: 1. DeLisser H, et al. 1994. Immunol. Today 15:490. 2. Newman P. 1997. J. Clin. Invest. 99:3. 3. Fawcett J, et al. 1995. J. Cell Biol. 128:1229. Application **Related Products: Product** Clone PE/Cy7 Mouse IgG1, κ Isotype Ctrl MOPC-21 FC, ICFC Cell Staining Buffer FC, ICC, ICFC RBC Lysis Buffer (10X) FC, ICFC Human TruStain FcX[™] (Fc Receptor Blocking Solution) FC, ICC, ICFC

For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.

*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.