


Anti-Mouse CD317 (BST2, PDCA-1) Alexa Fluor® 647

Catalog Number: 51-3171

Also Known As: plasmacytoid DC Ag-1, pDC Ag-1, BST 2, PDCA1

RUO: For Research Use Only



Staining of SJL splenocytes with Anti-Human/Mouse CD45R (B220) FITC (cat. 11-0452) (left) or Anti-Mouse CD11c PE (cat. 12-0114) (right) and Anti-Mouse CD317 (BST2, PDCA-1) Alexa Fluor® 647. Cells in the large forward scatter population were used for analysis.

Product Information

Contents: Anti-Mouse CD317 (BST2, PDCA-1) Alexa Fluor® 647

REF Catalog Number: 51-3171

Clone: eBio129c (129c)

Concentration: 0.2 mg/ml

Host/Isotype: Rat IgG2b

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

 Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

LOT Batch Code: Refer to Vial

 Use By: Refer to Vial

 Caution, contains Azide

Description

The eBio129c monoclonal antibody reacts with mouse PDCA-1 (BST2, CD317), a specific marker of plasmacytoid dendritic cells (pDC), also known as type I IFN-producing cells (IPC) in the naïve mouse. Mouse IPCs are typically CD11c+, CD11b-, B220+, Ly-6C+, and CD62L+. PDCA-1 is predominantly expressed by IPCs in the naïve mouse which represent a very minor population (<0.5%) of splenocytes. It is upregulated on numerous cell types following stimulation which triggers an IFN response. PDCA-1 cycles between cell surface and intracellular compartments and may function to regulate trafficking of secreted cytokines. PDCA-1 (BST2) is the protein recognized by the antibody 120G8.

The epitope recognized by eBio129c is distinct from eBio927; thus, the antibodies can be used to costain, purify and identify pDCs.

Applications Reported

This eBio129c (129c) antibody has been reported for use in flow cytometric analysis.

Applications Tested

This eBio129c (129c) antibody has been tested by flow cytometric analysis of SJL splenocytes. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

References

Blasius A.L., E. Giurisat, M. Celli, R.D. Schreiber, A.S. Shaw, M. Colonna. 2006. Bone Marrow Stromal Cell Antigen 2 Is a Specific Marker of Type I IFN-Producing Cells in the Naïve Mouse, but a Promiscuous Cell Surface Antigen Following IFN Stimulation. *J Immunol.* 177:3260-3265

Related Products

51-3172 Anti-Mouse CD317 (BST2, PDCA-1) Alexa Fluor® 647 (To Be Discontinued. Refer to Alternative Format: APC, cat. 17-3172) (eBio927)

51-4031 Rat IgG2b K Isotype Control Alexa Fluor® 647 (To Be Discontinued. Refer to NEW Format: eFluor® 660 cat. 50-4031)

Legal

Alexa Fluor® and Pacific Blue® are registered trademarks of and licensed under patents assigned to Molecular Probes, Inc. for research use only. This product is subject to an agreement between Molecular Probes, Inc. and eBioscience, and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications and corresponding foreign equivalents, owned by Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corp.). The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product for life science research or as an ASR. The buyer cannot use this product for manufacturing or for any other screening (specifically including use in combination with microarrays or High Content Screening) or testing purpose, other than as an ASR. For information on purchasing a license to this product for purposes other than life science research or use as an ASR, contact Molecular Probes, Inc.

Not for further distribution without written consent.

Copyright © 2000-2010 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com