


# Anti-Mouse/Rat Foxp3 Alexa Fluor® 647 (To Be Discontinued. Refer to Cat. No. 50-5773)

Catalog Number: 51-5773 Also Known As:Forkhead Box P3, Scurfin, JM2, Treg For Research Use Only. Not for use in diagnostic procedures.



**Product Information** 

**Contents:** Anti-Mouse/Rat Foxp3 Alexa Fluor® 647 (To Be Discontinued. Refer to Cat. No. 50-5773)

REF Catalog Number: 51-5773 Clone: FJK-16s Concentration: 0.2 mg/mL Host/Isotype: Rat IgG2a, kappa Intracellular staining of BALB/c splenocytes were Anti-Mouse CD4 PE (cat. 12-0042) and 0.5 ug of Anti-Mouse Foxp3 Alexa Fluor® 647 using the Foxp3 Staining Buffer Set (cat. 00-5523). Cells in the lymphocyte gate were used for analysis.

**Formulation:** aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

- **Temperature Limitation:** Store at 2-8°C. Do not freeze.
- Light sensitive material.
- LOT Batch Code: Refer to Vial
- Use By: Refer to Vial
- A Caution, contains Azide

## Description

The FJK-16s antibody reacts with mouse, rat, dog, porcine, bovine and cat Foxp3 also known as FORKHEAD BOX P3, SCURFIN, and JM2; cross reactivity of this antibody to other proteins has not been determined. Foxp3, a 49-55 kDa protein, is a member of the forkhead/winged-helix family of transcriptional regulators, and was identified as the gene defective in 'scurfy' (sf) mice. Constitutive high expression of foxP3 mRNA has been shown in CD4+CD25+ regulatory T cells (Treg cells), and ectopic expression of foxp3 in CD4+CD25- cells imparts a Treg phenotype in these cells.

Immunoblotting with FJK-16s antibody has mapped the epitope to amino acids 75-125 of the mouse Foxp3 protein. In the human, this region has been shown to be alternatively spliced at the mRNA level. Both the alternatively-spliced and non-spliced isoforms are present in the CD4+CD25+ subset of lymphocytes. Preliminary RT-PCR experiments have not revealed this alternatively-spliced isoform in mouse splenocytes, suggesting different gene regulation in the mouse and human.

## **Applications Reported**

This FJK-16s antibody has been reported for use in intracellular staining followed by flow cytometric analysis.

#### **Applications Tested**

This FJK-16s antibody has been tested by intracellular staining and flow cytometric analysis of mouse splenocytes using the Foxp3 Buffer Set (cat. 00-5523) and protocol. Please refer to Intracellular Staining Protocol on Best Protocol web page - Protocol B: One-step protocol for intracellular (nuclear) proteins. This antibody can be used at less than or equal to 0.25  $\mu$ g per test. A test is defined as the amount ( $\mu$ g) of antibody that will stain a cell sample in a final volume of 100  $\mu$ L. Cell number should be determined empirically but can range from 10<sup>5</sup> to 10<sup>8</sup> cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

#### References

Habicht A, Dada S, Jurewicz M, Fife BT, Yagita H, Azuma M, Sayegh MH, Guleria I. A link between PDL1 and T regulatory cells in fetomaternal tolerance. J Immunol. 2007 Oct 15;179(8):5211-9. (FJK-16s, IHC Frozen)

Heimesaat MM, Fischer A, Siegmund B, Kupz A, Niebergall J, Fuchs D, Jahn HK, Freudenberg M, Loddenkemper C, Batra A, Lehr HA, Liesenfeld O, Blaut M, Göbel UB, Schumann RR, Bereswill S. Shift towards pro-inflammatory intestinal bacteria aggravates acute murine colitis via Toll-like receptors 2 and 4. PLoS ONE. 2007 Jul 25;2(7):e662. (FJK-16s, IHC paraffin)

Leithäuser F, Meinhardt-Krajina T, Fink K, Wotschke B, Möller P, Reimann J. Foxp3-expressing CD103+ regulatory T cells accumulate

in dendritic cell aggregates of the colonic mucosa in murine transfer colitis. Am J Pathol. 2006 Jun;168(6):1898-909. (FJK-16s, IHC paraffin)

Kohm AP, McMahon JS, Podojil JR, Begolka WS, Degutes M, Kasprowicz DJ, Ziegler SF, Miller SD. Cutting Edge: Anti-CD25 Monoclonal Antibody Injection Results in the Functional Inactivation, Not Depletion, of CD4+CD25+ T Regulatory Cells. J Immunol. 2006 Mar 15;176(6):3301-5. (**FJK-16s**, IHC Frozen)

Suvas S, Azkur AK, Rouse BT.Qa-1b and CD94-NKG2a interaction regulate cytolytic activity of herpes simplex virus-specific memory CD8+ T cells in the latently infected trigeminal ganglia. J Immunol. 2006 Feb 1;176(3):1703-11. (FJK-16s, ICC)

Fontenot JD, Rasmussen JP, Williams LM, Dooley JL, Farr AG, Rudensky AY. Regulatory T cell lineage specification by the forkhead transcription factor foxp3. Immunity. 2005 Mar;22(3):329-41.

Hori S, Nomura T, Sakaguchi S. Control of regulatory T cell development by the transcription factor Foxp3. Science. 2003 Feb 14;299 (5609):1057-61.

## **Related Products**

00-5521 Foxp3 Fixation/Permeabilization Concentrate and Diluent 00-5523 Foxp3 / Transcription Factor Staining Buffer Set 12-0042 Anti-Mouse CD4 PE (RM4-5) 12-7979 Anti-Human/Mouse Foxp3 PE (Discontinued; Alternative clones: FJK-16s or PCH101) (eBio7979) 51-4321 Rat IgG2a K Isotype Control Alexa Fluor® 647 (eBR2a)

### Legal

Alexa Fluor® and Pacific Blue® are registered trademarks of and licensed under patents assigned to Molecular Probes, Inc. for research use only. This product is subject to an agreement between Molecular Probes, Inc. and eBioscience, and the manufacture, use, sale or import of this product may be subject to one or more U.S. patents, pending applications and corresponding foreign equivalents, owned by Molecular Probes, Inc. (a wholly owned subsidiary of Invitrogen Corp). The purchase of this product conveys to the buyer the non-transferable right to use the purchased amount of the product for life science research or as an ASR. The buyer cannot use this product for manufacturing or for any other screening (specifically including use in combination with microarrays or High Content Screening) or testing purpose, other than as an ASR. For information on purchasing a license to this product for purposes other than life science research or use as an ASR, contact Molecular Probes, Inc.

Not for further distribution without written consent. Copyright © 2000-2012 eBioscience, Inc. Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.eBioscience.com • info@eBioscience.com