

Product Data Sheet

PE/Cy5 anti-human CD33

Catalog # / Size: 303406 / 100 tests

Clone: WM53

Isotype: Mouse IgG1, κ

Workshop Number: IV M-505

Reactivity: Human, Cross-Reactivity: Chimpanzee

Preparation: The antibody was purified by affinity chromatography, and conjugated with

PE/Cy5 under optimal conditions. The solution is free of unconjugated


PE/Cy5 and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Human peripheral blood lymphocytes, monocytes and granulocytes stained with WM53 PE

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. Test

size products are transitioning from 20 μ l to 5 μ l per test. Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at

www.biolegend.com/testsize regarding the test size change.

Application Notes: Additional reported applications (for the relevant formats) include: immunoprecipitation, Westrn blotting³, and induction of cytokine production³. The LEAF™ purified antibody (Endotoxin <0.1 EU/μg, Azide-Free, 0.2 μm filtered) is

recommended for functional assays (Cat. No. 303410).

Cy3, Cy5, Cy5.5 and Cy7 are subject to proprietary rights of GE Healthcare Bio-Sciences Corp. and Carnegie Mellon University and made and sold under license from GE Healthcare Bio-Sciences Corp. Sale of this product is licensed

for research use only.

Application References: 1. Knapp W, et al. 1989. Leucocyte Typing IV. Oxford University Press. New York.

2. Favaloro E, et al. 1988. Br. J. Haematol. 69:163. 3. Garnache-Ottou F, et al. 2005. Blood 105:1256. (WB) 4. Chen YL, et al. 2012. PLoS One. 7:e47190. PubMed.

Description: CD33 is a 67 kD type I transmembrane glycoprotein also known as Siglec-3, gp67, and p67. It is a sialoadhesion

immunoglobulin superfamily member expressed on myeloid progenitors, monocytes, granulocytes, dendritic cells and mast cells. CD33 is absent on normal platelets, lymphocytes, erythrocytes and hematopoietic stem cells. CD33

HI98

functions as a sialic acid-dependent cell adhesion molecule with carbohydrate/lectin binding activity.

Antigen References: 1. Favaloro E, et al. 1988. Br. J. Haematol. 69:163.

PE anti-human CD15 (SSEA-1)

2. Freeman S, et al. 1995. Blood 85:2005.

Related Products: Product Clone Application

> FITC anti-human CD33 HIM3-4 PE anti-human CD14 M5E2 PE anti-human CD13 WM15 PE/Cy5 Mouse IgG1, κ Isotype Ctrl MOPC-21 FC, ICC, ICFC FC, ICFC Cell Staining Buffer

RBC Lysis Buffer (10X) Human TruStain FcX™ (Fc Receptor Blocking Solution) FC, ICC, ICFC



