

Product Data Sheet

Human peripheral blood leukocytes

stained with 581 Alexa Fluor® 647 and CD45 (HI30) FITC (top) or mIgG1, κ isotype control Alexa

Fluor® 647 and CD45 (HI30) FITC

(bottom). Cytograms were gated to display CD14 negative lymphocyte

CD45 FITC

population.

migG1, k Alexa

Alexa Fluor® 647 anti-human CD34

Catalog # / Size: 343507 / 25 tests

343508 / 100 tests

Clone: 581

Isotype: Mouse IgG1, κ

Workshop Number: V MA27

Reactivity: Human, Cross-Reactivity: Cynomolgus

Preparation: The antibody was purified by affinity chromatography, and conjugated with

Alexa Fluor® 647 under optimal conditions. The solution is free of

unconjugated Alexa Fluor® 647.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and

0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from

prolonged exposure to light. Do not freeze.



Applications: FC - Quality tested

IF, IHC - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent

staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is 5 μ I per million cells or 5 μ I per 100 μ I of whole blood. It is recommended that the reagent be titrated for optimal

performance for each application.

* Alexa Fluor® 647 has a maximum emission of 668 nm when it is excited at

633nm / 635nm.

Alexa Fluor® 647 is a registered trademark of Molecular Probes, Inc. Alexa Fluor® 647 dye antibody conjugates are sold under license from Molecular Probes, Inc. for research use only, except for use in combination with microarrays and high content screening, and are covered by pending and

issued patents.

Application Notes: The 581 antibody recognizes the class III group epitope which is resistant to

sialidase/glycolyprotease and chymopapain treatment. Additional reported applications (for the relevant formats) include: immunohistochemical staining of paraffin-embedded tissue sections⁵ and immunofluorescence⁶.

Application References: 1. Schlossman SF, et al. 1995. Leukocyte Typing V:White Cell Differentiation Antigen. New York:Oxford University Press.

2. Felschow DM, et al. 2001. Blood 97:3768

3. Rudin CE, et al. 1997. Br. J. Haematol. 97:488.

Yoshino N, et al. 2000. Exp. Anim. (Tokyo) 49:97. (FC)

5. Skowasch D, et al. 2003. Cardiovasc Rés. 60:684. (IHC) 6. Umland O, et al. 2003. J. Histochem. Cytochem. 51:977. (IF)

Description: CD34, also known as gp105-120, is a type I monomeric sialomucin-like glycophosphoprotein with an approximate molecular weight of 105-120 kD. Selectively expressed on the majority of hematopoietic stem/progenitor cells, bone

marrow stromal cells, capillary endothelial cells, embryonic fibroblasts, and some nervous tissue, CD34 is a commonly used marker to identify human hematopoietic stem/progenitor cells. According to the differential sensitivity to enzymatic cleavage, four groups of epitopes of CD34 have been described. CD34 mediates cell adhesion and

lymphocytes homing through binding to L-selectin and E-selectin ligands.

Antigen References: 1. Krause DS, et al. 1996. Blood 87:1.

2. Puri KD, et al. 1995. J. Cell Biol. 131:261.

3. Zola H, et al. 2007. Leukocyte and Stromal Cell Molecules: The CD Markers. John Wiley & Sons Inc, Hoboken New

Jersev.

Related Products: Product

Alexa Fluor® 647 Mouse IgG1, κ Isotype Ctrl (FC)

Cell Staining Buffer


RBC Lysis Buffer (10X)

Clone MOPC-21 Application FC, IF FC, ICC, ICFC FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.





BioLegend Inc., 9727 Pacific Heights Blvd, San Diego, CA 92121 www.biolegend.com Toll-Free Phone: 1-877-Bio-Legend (246-5343) Phone: (858) 768-5800 Fax: (877) 455-9587



