

## FITC anti-human CD40

**Catalog # / Size:** 313003 / 25 tests  
313004 / 100 tests

**Clone:** HB14

**Isotype:** Mouse IgG1,  $\kappa$


**Workshop Number:** V CD40.5

**Reactivity:** Human, **Cross-Reactivity\*:** Rhesus

**Preparation:** The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

**Storage:** The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with HB14 FITC

## Applications:

**Applications:** FC - Quality tested

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20  $\mu$ l to 5  $\mu$ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100  $\mu$ l staining volume or per 100  $\mu$ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at [www.biolegend.com/testsize](http://www.biolegend.com/testsize) regarding the test size change.

**Application Notes:** Additional reported applications (for the relevant formats) include: costimulation of B cell proliferation, partial inhibition of CD40 binding to CD40L, and prevention of B cell apoptosis.<sup>1</sup> Alone, or in combination with TLR ligands, clone HB14 stimulates B cells to produce IL-10 and differentiates it into regulatory B10 (IL-10 producing B cells).<sup>7</sup> The LEAF™ purified antibody (Endotoxin <0.1 EU/ $\mu$ g, Azide-Free, 0.2  $\mu$ m filtered) is recommended for functional assays (Cat. No. 313010).

**Application References:**

1. Pound JD, *et al.* 1999. *Int. Immunol.* 11:11. (Costim)
2. Schlossman S, *et al.* Eds. 1995. *Leucocyte Typing V.* Oxford University Press. New York.
3. Armengol MP, *et al.* 2001. *Am. J. Pathol.* 159:861.
4. Cavanagh LL, *et al.* 2005. *Arthritis Res. Ther.* 7:R230.
5. Jayakumar A, *et al.* 2008. *Infect Immun.* 76:2138. PubMed
6. Sestak K, *et al.* 2007. *Vet. Immunol. Immunopathol.* 119:21.
7. Iwata Y, *et al.* 2011. *Blood.* 117:530. PubMed

**Description:** CD40 is a 48 kD type I glycoprotein also known as BP50. It is a member of the TNFR superfamily primarily expressed on B cells, macrophages, follicular dendritic cells, endothelial cells, fibroblasts, and at low levels on plasma cells. CD40 has been reported to be involved in B cell differentiation, costimulation, isotype class-switching, and protection of B cells from apoptosis. Additionally, CD40 is important for T cell-B cell interactions. The ligand of CD40 is CD154 (CD40 ligand). The HB14 antibody has been reported to promote B cell proliferation in the presence of anti-IgM, IL-4 or PMA, partially block CD40 binding to CD40L and rescue B cells from apoptosis.

**Antigen References:**

1. Banchereau J, *et al.* 1994. *Annu. Rev. Immunol.* 12:881.
2. Foy T, *et al.* 1996. *Annu. Rev. Immunol.* 14:591.

### Related Products: Product

FITC Mouse IgG1, $\kappa$ Isotype Ctrl	MOPC-21
Cell Staining Buffer	
RBC Lysis Buffer (10X)	
FITC anti-human CD154	24-31
Human TruStain FcX™ (Fc Receptor Blocking Solution)	

**Clone**  
MOPC-21

24-31

### Application

FC, ICFC  
FC, ICC, ICFC  
FC, ICFC  
FC  
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.