


Anti-Mouse CD16/CD32 eFluor® 450

Catalog Number: 48-0161

Also known as: FCGR3, IGFR3; FCGR2, IGFR2; FC Receptor Block

RUO: For Research Use Only. Not for use in diagnostic procedures.



Staining of C57BL/6 splenocytes with 0.25 ug of Rat IgG2a K Isotype Control eFluor® 450 (cat. 48-4321) (blue histogram) or 0.25 ug of Anti-Mouse CD16/CD32 eFluor® 450 (purple histogram). Total viable cells were used for analysis.

Product Information

Contents: Anti-Mouse CD16/CD32 eFluor® 450

Catalog Number: 48-0161

Clone: 93

Concentration: 0.2 mg/mL

Host/Isotype: Rat IgG2a, lambda

Formulation: aqueous buffer, 0.09% sodium azide, may contain carrier protein/stabilizer

Temperature Limitation: Store at 2-8°C. Do not freeze. Light sensitive material.

Batch Code: Refer to vial

Use By: Refer to vial

Caution, contains Azide

Description

The 93 monoclonal antibody reacts with an epitope shared by mouse CD16 and CD32. CD16 (Fc gamma III Receptor) and CD32 (Fc gamma II Receptor) are the low affinity receptors for the mouse IgG Fc portion and are expressed by B cells, monocyte/macrophages, NK cells, and neutrophils.

Applications Reported

This 93 antibody has been reported for use in flow cytometric analysis.

eFluor® 450 is a replacement for Pacific Blue®. eFluor® 450 emits at 456 nm and is excited with the Violet laser (405 nm). Please make sure that your instrument is capable of detecting this fluorochrome.

Applications Tested

This 93 antibody has been tested by flow cytometric analysis of mouse splenocytes. This can be used at less than or equal to 0.5 µg per test. A test is defined as the amount (µg) of antibody that will stain a cell sample in a final volume of 100 µL. Cell number should be determined empirically but can range from 10⁵ to 10⁸ cells/test. It is recommended that the antibody be carefully titrated for optimal performance in the assay of interest.

eFluor™ 450 is a replacement for Pacific Blue®. eFluor™ 450 emits at 456 nm and is excited with the Violet laser (405 nm). Please make sure that your instrument is capable of detecting this fluorochrome.

References

Oliver, A. M., J. C. Grimaldi, et al. Independently ligating CD38 and Fc gammaRIIB relays a dominant negative signal to B cells. 1999. Hybridoma 18(2): 113-9.

Related Products

48-4321 Rat IgG2a K Isotype Control eFluor® 450 (eBR2a)

Not for further distribution without written consent.

Copyright © 2000-2012 eBioscience, Inc.

Tel: 888.999.1371 or 858.642.2058 • Fax: 858.642.2046 • www.ebioscience.com • info@ebioscience.com