

## Purified anti-human CD59

**Catalog # / Size:** 304701 / 25 µg  
304702 / 100 µg

**Clone:** p282 (H19)

**Isotype:** Mouse IgG2a, κ

**Workshop Number:** V S006


**Reactivity:** Human, **Cross-Reactivity\*:** Chimpanzee, Baboon, Cynomolgus, Rhesus

**Preparation:** The antibody was purified by affinity chromatography.

**Formulation:** Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

**Concentration:** 0.5 mg/ml

**Storage:** The antibody solution should be stored undiluted at 4°C.



Human peripheral blood lymphocytes stained with purified p282, followed by anti-mouse IgGs FITC

## Applications:

**Applications:** FC - *Quality tested*

**Recommended Usage:** Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is  $\leq 2.0$  µg per  $10^6$  cells in 100 µl volume or 100 µl of whole blood. It is recommended that the reagent be titrated for optimal performance for each application.

**Application References:** 1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.

**Description:** CD59 is a 19-25 kD glycosylphosphatidylinositol (GPI)-anchored glycoprotein also known as protectin, MACIF, and H19. It is broadly expressed on hematopoietic and non-hematopoietic cells. CD59 inhibits the cytolytic activity of complement by binding to C9, inhibiting incorporation into C5b-8, and preventing the terminal steps in complement polymerization of the membrane attack complex. CD59 has also been reported to play a role in T cell activation.

**Antigen References:** 1. Davies A, *et al.* 1993. *Immunol. Res.* 12:258.  
2. Lachmann P. 1991. *Immunol. Today* 12:312.  
3. Liszewski M, *et al.* 1996. *Adv. Immunol.* 61:201.

### Related Products:

Product	Clone	Application
APC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Biotin Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC, ELISA, IHC, IF, WB
FITC Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Purified Mouse IgG2a, κ Isotype Ctrl	MOPC-173	FC, ICC, IF, IHC, IP, WB
PE Goat anti-mouse IgG (minimal x-reactivity)	Poly4053	FC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
Purified anti-human CD55	JS11	FC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



\*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, [www.biolegend.com/ordering#license](http://www.biolegend.com/ordering#license)). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.