

Product Data Sheet

FITC anti-human CD59

Catalog # / Size: 304706 / 100 tests

Clone: p282 (H19)

Isotype: Mouse IgG2a, κ


Workshop Number: V S006

Reactivity: Human, **Cross-Reactivity*:** Chimpanzee, Baboon, Cynomolgus, Rhesus

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with p282(H19) FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References: 1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.

Description: CD59 is a 19-25 kD glycosylphosphatidylinositol (GPI)-anchored glycoprotein also known as protectin, MACIF, and H19. It is broadly expressed on hematopoietic and non-hematopoietic cells. CD59 inhibits the cytolytic activity of complement by binding to C9, inhibiting incorporation into C5b-8, and preventing the terminal steps in complement polymerization of the membrane attack complex. CD59 has also been reported to play a role in T cell activation.

Antigen References: 1. Davies A, *et al.* 1993. *Immunol. Res.* 12:258.
2. Lachmann P. 1991. *Immunol. Today* 12:312.
3. Liszewski M, *et al.* 1996. *Adv. Immunol.* 61:201.

Related Products:

Product
FITC Mouse IgG2a, κ Isotype Ctrl

Cell Staining Buffer

RBC Lysis Buffer (10X)

FITC anti-human CD55

Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone

MOPC-173

JS11

Application

FC, ICFC

FC, ICC, ICFC

FC, ICFC

FC

FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.