

FITC anti-human CD55

Catalog # / Size: 311305 / 25 tests
311306 / 100 tests

Clone: JS11

Isotype: Mouse IgG1, κ


Workshop Number: VI N-L060

Reactivity: Human

Preparation: The antibody was purified by affinity chromatography, and conjugated with FITC under optimal conditions. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood lymphocytes stained with JS11 FITC

Applications:

Applications: FC - Quality tested

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application References:

1. Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
2. Peyron P, *et al.* 2000. *J. Immunol.* 165:5186
3. Nevo Y, *et al.* 2013. *Blood.* 121:129. PubMed.

Description: CD55 is a 60-70 kD glycosylphosphatidylinositol (GPI)-anchored single chain glycoprotein also known as decay-accelerating factor (DAF). It is expressed on hematopoietic cells including erythrocytes and many non-hematopoietic cells. CD55 accelerates the dissociation of the components of the C3-convertases (namely C2a from C4b in the C4bC2a complex, a C3-convertase of the classical pathway, and factor Bb from the C3bBb complex, a C3-convertase of the alternative pathway) to protect cells from inappropriate damage caused by autologous complement. CD55 has been reported to reduce the efficiency of NK cell lysis and induce signal transduction in T cells. CD55 has also been shown to interact with CD97 and bind to Coxsackie and Echovirus.

Antigen References:

1. Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York.
2. Hamann J, *et al.* 1996. *J. Exp. Med.* 184:1185.
3. Fujita T, *et al.* 1987. *J. Exp. Med.* 166:1221

Related Products:

Product	Clone	Application
FITC anti-human CD55	p282 (H19)	FC
FITC Mouse IgG1, κ Isotype Ctrl	MOPC-21	FC, ICFC
Cell Staining Buffer		FC, ICC, ICFC
RBC Lysis Buffer (10X)		FC, ICFC
Human TruStain FcX™ (Fc Receptor Blocking Solution)		FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.