

FITC anti-human CD57

Catalog # / Size: 322305 / 25 tests
322306 / 100 tests

Clone: HCD57

Isotype: Mouse IgM, κ

Reactivity: Human

Preparation: The antibody was conjugated with FITC under optimal conditions, and is at >85% purity. The solution is free of unconjugated FITC.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**

Applications:

Applications: FC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: The HCD57 antibody has been shown to recognize human CD57 and to be useful for flow cytometry.

Application References: 1. Kaplan RC, *et al.* 2011. *J. Infect Dis.* 10:76. PubMed
2. Lutz CT, *et al.* 2011. *J. Immunol.* 186:4590. PubMed

Description: CD57, also known as HNK-1, NK-1, and Leu-7, is a 100-115 kD oligosaccharide antigenic determinant expressed on a variety of proteins, lipids, and chondroitin sulfate proteoglycans. CD57 is expressed on a subset of peripheral blood lymphocytes, including NK cells and CD8⁺ T cells, and is also expressed on neural cells and striated muscle. CD57 is not expressed on red cells, granulocytes, monocytes, or platelets. While the function of CD57 is unknown, binding to L-selectin, P-selectin, and a fragment of laminin suggests that CD57 may be involved in cell-matrix interactions. CD57 is increased in some disease states associated with CD4/CD8 imbalances (AIDS, autoimmune disease, viral infections, and allograft transplants).

Antigen References: 1. Schubert J, *et al.* 1989. In *Leucocyte Typing IV* (Knapp W, ed) Oxford University Press Oxford pp 711-714.
2. Palmer BE, *et al.* 2005. *J. Immunol.* 175:8415.
3. Schachner M, *et al.* 1995. *Trends Neurosci.* 18:183.
4. Wood KL, *et al.* 2005. *Clin. Immunol.* 117:294.

Related Products:


Product
Cell Staining Buffer
FITC Mouse IgM, κ Isotype Ctrl
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone


MM-30

Application

FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



Human peripheral blood lymphocytes were stained with CD8 APC and CD57 (clone HCD57) FITC (top) or mouse IgM FITC isotype control (bottom).



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.