

Purified anti-human CD51

Catalog # / Size: 327901 / 25 µg
327902 / 100 µg

Clone: NKI-M9

Isotype: Mouse IgG2a, κ

Workshop Number: IV 103

Immunogen: Melonama cells


Reactivity: Human

Preparation: The antibody was purified by affinity chromatography.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide.

Concentration: 0.5 mg/ml

Storage: The antibody solution should be stored undiluted at 4°C.



Human melanoma cell line M21 stained with purified NKI-M9, followed by anti-mouse IgG FITC

Applications:

Applications: FC - Quality tested
IP, ELISA - Reported in the literature

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. For immunofluorescent staining, the suggested use of this reagent is ≤2.0 µg per million cells in 100 µl volume. It is recommended that the reagent be titrated for optimal performance for each application.

Application References:

- Knapp W, et al. Eds. 1989. Leucocyte Typing IV. Oxford University Press. New York.
- Defilippi P, et al. 1991. *J. Cell Biol.* 114:855. PubMed
- Burdick MM, et al. 2003. *Am. J. Physiol. Cell Ph.* 284:C977. PubMed
- Grzeszkiewicz TM, et al. 2001. *J. Biol. Chem.* 276:21943.
- Sonnenberg A, et al. 1990. *J. Cell Biol.* 110:2145. PubMed

Description: CD51 is a type I integral membrane glycoprotein, known as vitronectin receptor α chain, or integrin α_v. It forms heterodimer with integrin β1 (CD29), β3 (CD61), β5, β6, or β8. CD51 contains two disulfide-linked subunits of 125 kD and 24 kD, and is expressed on endothelial cells, fibroblasts, macrophages, platelets, osteoclasts, neuroblastoma, melanoma, and hepatoma cells. Many extracellular matrix proteins with RGD-motifs are CD51 ligands. In association with its β chains, CD51 binds vitronectin, von Willebrand factor, fibronectin, thrombospondin, osteopontin, fibrinogen, and laminin. CD51, as an adhesion molecule, plays important roles in leukocytes homing and rolling, mediates bone absorption and angiogenesis.

Antigen References:

- Nesbitt S, et al. 1993. *J. Biol. Chem.* 268:16737.
- Zola H, et al. 2007. Leukocyte and Stromal Cell Molecules: The CD Markers Wiley-Liss A John Wiley & Sons Inc, Publication

Related Products:

Product
Purified Mouse IgG2a, κ Isotype Ctrl
Cell Staining Buffer
RBC Lysis Buffer (10X)

Clone
MOPC-173

Application
FC, ICC, IF, IHC, IP, WB
FC, ICC, ICFC
FC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.