

APC anti-human CD64

Catalog # / Size: 305013 / 25 tests
305014 / 100 tests

Clone: 10.1

Isotype: Mouse IgG1, κ

Workshop Number: VI MA36


Immunogen: Human rheumatoid synovial fluid cells and fibronectin-purified monocytes.

Reactivity: Human, **Cross-Reactivity:** Chimpanzee, Baboon, Cynomolgus, Rhesus, Capuchin Monkey, Squirrel Monkey

Preparation: The antibody was conjugated with APC under optimal conditions, and is at >85% purity. The solution is free of unconjugated APC and unconjugated antibody.

Formulation: Phosphate-buffered solution, pH 7.2, containing 0.09% sodium azide and 0.2% (w/v) BSA (origin USA).

Storage: The antibody solution should be stored undiluted at 4°C and protected from prolonged exposure to light. **Do not freeze.**



Human peripheral blood monocytes stained with 10.1 APC

Applications:

Applications: FC - *Quality tested*

Recommended Usage: Each lot of this antibody is quality control tested by immunofluorescent staining with flow cytometric analysis. **Test size products are transitioning from 20 μ l to 5 μ l per test.** Please check your vial or your CoA to find the suggested use of this reagent per million cells in 100 μ l staining volume or per 100 μ l of whole blood. It is recommended that the reagent be titrated for optimal performance for each application. Read more at www.biolegend.com/testsize regarding the test size change.

Application Notes: Clone 10.1 recognizes the EC3 epitope of CD64. Additional reported applications (for the relevant formats) include: blocking of human IgG3 and murine IgG2a binding to Fc γ RI^{2,5,6,11} and immunohistochemical staining of acetone-fixed frozen tissue sections.

- Application References:**
- McMichael A, *et al.* Eds. 1987. Leucocyte Typing III. Oxford University Press. New York.
 - Schlossman S, *et al.* Eds. 1995. Leucocyte Typing V. Oxford University Press. New York. p. 874.
 - Kishimoto T, *et al.* Eds. 1997. Leucocyte Typing VI. Garland Publishing Inc. London.
 - Höll V, *et al.* 2004. *J. Immunol.* 173:6274.
 - Hober D, *et al.* 2002. *J. Gen. Virol.* 83:2169.
 - Cho HJ, *et al.* 2007. *Physiol Genomics* 149:60.
 - van Tits L, *et al.* 2005. *Arterioscler Thromb Vasc Biol.* 25:717. PubMed
 - Bruhns P, *et al.* 2008. *Blood* 113:3716. PubMed
 - Yoshino N, *et al.* 2000. *Exp. Anim. (Tokyo)* 49:97. (FC)
 - Carter DL, *et al.* 1999. *Cytometry* 37:41. (FC)
 - Dougherty GJ, *et al.* 1987. *Eur. J. Immunol.* 17:1453.
 - Shapiro H, *et al.* 2013. *J Clin Endocrinol Metab.* 98:1173. PubMed.

Description: CD64 is a 72 kD single chain type I glycoprotein also known as Fc γ RI and FcR I. CD64 is a member of the immunoglobulin superfamily and is expressed on monocytes/macrophages, dendritic cells, and activated granulocytes. The expression can be upregulated by IFN- γ stimulation. CD64 binds IgG immune complex. It plays a role in antigen capture, phagocytosis of IgG/antigen complexes, and antibody-dependent cellular cytotoxicity (ADCC).

- Antigen References:**
- Hulett M, *et al.* 1994. *Adv. Immunol.* 57:1.
 - van de Winkel J, *et al.* 1993. *Immunol. Today* 14:215.

Related Products:

Product
APC Mouse IgG1, κ Isotype Ctrl (FC)
Cell Staining Buffer
RBC Lysis Buffer (10X)
Human TruStain FcX™ (Fc Receptor Blocking Solution)

Clone
MOPC-21

Application
FC
FC, ICC, ICFC
FC, ICFC
FC, ICC, ICFC



For research use only. Not for diagnostic use. Not for resale. BioLegend will not be held responsible for patent infringement or other violations that may occur with the use of our products.



*These products may be covered by one or more Limited Use Label Licenses (see the BioLegend Catalog or our website, www.biolegend.com/ordering#license). BioLegend products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products, reverse engineer functionally similar materials, or to provide a service to third parties without written approval of BioLegend. By use of these products you accept the terms and conditions of all applicable Limited Use Label Licenses. Unless otherwise indicated, these products are for research use only and are not intended for human or animal diagnostic, therapeutic or commercial use.